



## Hematological and Oxidative Stress Markers Analysis for Detection and Prediction of Osteoporosis in Post-menopausal Women

Derouiche Samir<sup>1,2\*</sup>, Haddig Nour El-houda<sup>1</sup>, Zerzour Aicha<sup>1</sup>

<sup>1</sup>Department of Cellular and Molecular Biology, Faculty of Natural Sciences and Life, University of El Oued, El-Oued 39000, Algeria

<sup>2</sup>Laboratory of Biodiversity and Application of Biotechnology in the Agricultural Field, University of El Oued, El-Oued 39000, Algeria.

### ABSTRACT

The main purpose of this work was to analyze some biological and oxidative stress markers to predict and diagnose osteoporosis in postmenopausal women. For the experiment, we have chosen 20 healthy menopausal women as control and 20 menopausal women who had osteoporosis. Some biochemical, hematological, and oxidative stress parameters were measured. The sensitivity and specificity of oxidative stress biomarkers in serum, erythrocytes, and leucocytes were estimated using Receiver Operating Characteristics (ROC) curve design. The results of the study demonstrate a significant change ( $P < 0.05$ ) in biochemical and hematological markers during osteoporosis in post-menopausal women. In addition, results show a significant enhancement of MDA content and reduction in GSH, SOD, Catalase, and TAC levels in patients compared to control. Some oxidative stress markers identify as risk factors ( $P < 0.05$ ,  $OR > 1$ ) and represented an important significant specificity ( $P < 0.05$ ) for osteoporosis diagnosis such as serum MDA (AUC=99.6%), erythrocytic GSH (AUC=100%), and leukocytic SOD (AUC=62.5%), with a strong correlation ( $P < 0.05$ ) between GSH levels and serum calcium and ALP changes. In conclusion, several clinical factors contributed to the evolution of osteoporosis in postmenopausal women. Oxidative stress and hematologic markers represent very important diagnostic and predictive factors for osteoporosis in women during the post-menopause period.

**Keywords:** Osteoporosis, Post-menopause, Risk factors, Oxidative stress

**Corresponding author:** Derouiche Samir

**e-mail** ✉ [dersamebio@gmail.com](mailto:dersamebio@gmail.com)

**Received:** 29 March 2021

**Accepted:** 27 June 2021

### INTRODUCTION

Menopause implies the continual cessation of the monthly cycle and the conclusion of regenerative ability (Weiss *et al.*, 2004). The menopausal evolution includes a period of dynamic alterations in non-reproductive and reproductive tissues. This evolution is considered to have a main effect on the etiology of symptoms including night sweats, uterine bleeding problems, vulvovaginal atrophy, hot flashes, and mood changes (Zeidabadi *et al.*, 2020). Osteoporosis is defined as an illness that is specified by disruption of bone microarchitecture, deterioration of bone tissue, and low bone mass: it may result in compromised bone quality and an increment in the chance of breaks (Al-Shali *et al.*, 2019; Nambi *et al.*, 2020; Zerzour *et al.*, 2020). The social and economic burden of osteoporosis is enhancing continuously due to the world population's aging (Sözen *et al.*, 2017). Nowadays, osteoporosis is projected to affect about 14 million adults above 50 years old by the year 2020, affecting above 10 million people in the United States; (Wright *et al.*, 2014). Several late landmarks in scientific studies have indicated that oxidative stress is a significant factor in human beings, resulting in physiological and metabolic modifications and different illnesses in the body (Atoussi *et al.*, 2021). Oxidative stress is an unusual condition

as a result of an overabundance generation of oxidants in comparison to the antioxidants (Derouiche *et al.*, 2017) that has been regarded as the major cause of many pathologies (Chetehouna *et al.*, 2020). Minerals have a significant effect on the blood clotting, receiving and sending signals, formation of bones, production of cell energy, oxygen transportation, keeping normal heartbeat, synthesizing and metabolizing proteins and fats, providing immunity to the body, acting as coenzymes, and helping nervous system work properly (Tomlinson *et al.*, 2020). Considering this information, the purpose of our work is based on the determination of the variation and specificity of some oxidative and hematological stress markers in the prediction and diagnosis following up of osteoporosis in menopausal women.

### MATERIALS AND METHODS

#### Subjects and Study Design

Ethical approval was obtained from the ethics committee (28 EC/DCMB/FNSL/EU2020) of the department of cellular and molecular biology, Faculty of Natural Sciences and Life, El-Oued University. This work was applied to 40 volunteer women aged between 43-54 years, they were divided into two groups; a group of 20 healthy control women with an average age of  $48.26 \pm 0.43$  years old, a group of 20 menopause women having osteoporosis with the mean age  $48.77 \pm 0.21$  years old. In addition, in this study, we included voluntary women living in the Guemar El-Oued region, who were women suffering

from osteoporosis aged from 43 to 54 and like control women were in good health and did not have any pathology. Moreover, in the present research, we excluded all women who were suffering from other acute or chronic pathology and all women using drugs for 30 days or during post-menopausal periods.

#### Sample Collection and Analyses

Blood sampling for both groups was done during morning fasting. After the blood sampling, the blood was collected in two types of tubes, in the anticoagulant tube (EDTA), for hematological and oxidative stress (MDA, GSH, SOD, and CAT) parameters assays. In dry tubes, samples were centrifuged at 3000 rpm for 10 minutes, and then the serum was recovered to achieve the dosage of biochemistry parameters: Glucose, calcium, iron, PAL, vitamin C, and total antioxidant ORAC. Serum glucose, calcium, iron, and PAL were determined by the Semi-auto Analyzer Mindray BA-88A and measured using commercial kits from Biomaghreb, (Biomaghreb: glucose-20121, calcium-20051, iron-20064, and PAL-20015. Hematological analysis (FNS) was performed by the hematology Auto analyzer (Mindray).

#### Oxidative Markers Measurement

MDA was measured using TBA reagent according to the method described by Sastre *et al.* (2000). The level of reduced Glutathione is determined according to the Weak and Cory (1988). The catalase activity consisting measuring the catalase-induced loss of H<sub>2</sub>O<sub>2</sub> contained in the sample according to the

method of Aebi (1984). The assay method of SOD activity was the method of Beauchamp and Fridovich, (1971). The plasma vitamin C is measured according to the method of Jagota and Dani (1982) using the Folin reagent. The total antioxidant power of the serum and its capacity to absorb free oxygen radicals (ORAC: Oxygen Radical Absorbance Capacity) were evaluated according to the method of Oyaizu, (1986).

#### Statistical Analysis

Statistical analysis was performed by the SPSSV20.0 software. Results comparisons were carried out by using the Student T-test to contrast means among the groups. Correlation analysis was carried out using the Pearson Correlation test and regression analysis was used for other analyses and statistical data. Differences were considered statically significant at p <0.05.

## RESULTS AND DISCUSSION

#### Description of the Study Population

The general data of socioeconomic characteristics of the two groups of subjects include age, Body Weight, BMI, social case, number of children, job, educational level, and Blood group. Most of these indicators do not have any statistically significant (P > 0.05) differences but serum calcium was decreased in the patients' group compared to the control. Other baseline characteristics between the two main groups are summarized in **Table 1**.

**Table 1.** Demographic, Clinical and Laboratory Features between the Study Groups

Parameter	Control (n=20)	Patients (n=20)	P- value	
Age (ys)	48.26±0.43	48.77±0.21	0.114	
Body Weight (kg)	68.60±0.243	71.40±0.61	0.080	
Height (cm)	162.76±0.164	161.98±0.130	0.204	
Body Mass index	26.01±0.182	27.39±0.108	0.152	
Blood glucose (g/l)	0.82±0.023	0.92±0.018	0.009	
Serum Calcium (g/l)	74.31±1.08	65.15±2.04	0.006	
Serum ALP (U/l)	129.0±10.2	182±13.9	0.002	
Number of children	5,333±0,18	6,06±0,18	0,081	
Social Case (%)	Married	81	96	0.014
	Single	7	2	0.024
	Divorced	10	1	0.000
	Widow	2	1	0.072
Job (%)	Worker	25	10	0.000
	Housewife	75	90	0.000
Educational Level (%)	Illiterate	13	14	0.062
	Primary	15	28	0.002
	Junior high	26	24	0.123
	High School	26	23	0.092
	Higher education	20	11	0.014
Blood Group (%)	A	43.33	20	0.000
	B	10	10	0.231
	AB	6.67	13	0.002
	O	40	57	0.051

#### Hematological Parameters

The illustrated results of the hematological parameters in (**Table 2**) show that in the osteoporosis patients group there

was a significant increase (P<0.05) in WBC, Lymphocytes, RBC, and in HCT and PLT (P0.001>) levels and no significant change of hemoglobin level in comparison with the control group.

**Table 2.** Changes in the Haematological Levels in Control and Osteoporosis Patients

Parameter	Control (n=20)	patients (n=20)	P-value
White Blood Cells ( $\times 10^3/\mu\text{l}$ )	5,669 $\pm$ 0,286	7,169 $\pm$ 0,56	0,020
Lymphocytes ( $\times 10^3/\mu$ )	2,042 $\pm$ 0,073	2,200 $\pm$ 0,195	0,432
Hemoglobin (g/dl)	125,73 $\pm$ 2,49	130,06 $\pm$ 1,83	0,032
Red Blood Cells ( $\times 10^6/\mu\text{l}$ )	4,562 $\pm$ 0,055	4,752 $\pm$ 0,086	0,043
Hematocrite (%)	39,8 $\pm$ 0,719	43,256 $\pm$ 0,603	0,000
Platelets ( $\times 10^3/\mu\text{l}$ )	136,06 $\pm$ 1,86	255,6 $\pm$ 16,4	0,000

**Oxidative Stress Markers**

The analysis of blood oxidative stress parameters in control and patient are shown in **Table 3**, The results show a significant decrease of GSH level in erythrocyte (P<0.001), leukocyte (P<0.001) and serum (P<0.001) and also decrease of

catalase and SOD activities in leukocyte (P<0.001) and in serum (P<0.001, P<0.05) and TAC activity in serum and a significant increase of MDA level in leukocyte (P<0.05) and serum (P<0.001) in osteoporosis patients group in comparison to the control group.

**Table 3.** Parameters of Oxidative Stress in Blood of Control and Osteoporosis Patients

Parameter	Control (n=20)	Patient (n=20)	P-value	
MDA (nM/mg of Hb)	Erythrocytes	9,06 $\pm$ 1,03	7,587 $\pm$ 0,780	0,083
	Leukocytes	7,566 $\pm$ 0,785	12,11 $\pm$ 1,76	0,021
	Serum	28,02 $\pm$ 2,84	92,5 $\pm$ 10,4	0,000
GSH (nM/mg of Hb)	Erythrocytes	1,065 $\pm$ 0,041	0,898 $\pm$ 0,027	0,000
	Leukocytes	1,448 $\pm$ 0,217	0,765 $\pm$ 0,107	0,000
	Serum	3,751 $\pm$ 0,146	2,969 $\pm$ 0,295	0,021
CAT (UI/g of Hb)	Leukocytes	0,032 $\pm$ 0,003	0,021 $\pm$ 0,001	0,000
	Serum	0,11 $\pm$ 0,006	0,099 $\pm$ 0,004	0,031
SOD (U/mg of Hb)	Leukocytes	15,250 $\pm$ 0,423	12,929 $\pm$ 0,339	0,000
	Serum	78,56 $\pm$ 1,99	73,43 $\pm$ 1,03	0,000
Vit C ( $\mu\text{g/ml}$ )	Serum	61,63 $\pm$ 3,47	66,38 $\pm$ 3,86	0,238
TAC (U/l)	Serum	21,24 $\pm$ 1,55	14,88 $\pm$ 1,69	0,002

MDA=Malondialdehyde; GSH=Glutathione; CAT= Catalase SOD=Superoxide dismutase; TAC=Total antioxidant capacity; Vit C =Vitamin C.

**Study of Odds Ratio Values of Biochemical and Oxidative Stress Markers**

Odds Ratio (OR) values for some oxidative stress parameters and biochemical markers of controls and patients groups (**Table 4**) show that decreased serum ORAC, leukocyte SOD, erythrocyte GSH, leukocyte GSH, leukocyte catalase, and serum

calcium are shown to be important risk factors for osteoporosis OR (6.303-106.375) with P<0.05. Reciprocally, decreased serum MDA and serum PAL are protective factors against osteoporosis in the study population (OR=0.032; P=0.000, OR=0.084; P=0.000), respectively.

**Table 4.** Odds Ratio Value of Biochemical and Oxidative Stress Markers

	Control %	Patient %	OR	CI95%	P-value
<b>Serum ORAC</b>			6.303	2.604-15.255	0.000
Positive	32	11			
Negative	18	39			
<b>Leukocyte SOD</b>			106.375	13.475-839.745	0.000
Positive	37	01			
Negative	16	46			
<b>Erythrocyte GSH</b>			37.161	4.796-287.936	0.000
Positive	32	01			
Negative	31	36			
<b>Leukocyte GSH</b>			18.951	2.428-147.909	0.000
Positive	21	01			
Negative	41	37			
<b>Leukocyte catalase</b>			8.957	2.424-33.103	0.000
Positive	17	3			
Negative	31	49			

<b>Serum MDA</b>			0.032	0.004-0.248	0.000
Positive	30	50			
Negative	19	1			
<b>Serum calcium</b>			18.222	5.035-65.946	0.000
Positive	32	3			
Negative	24	41			
<b>Serum PAL</b>			0.084	0.030-0.233	0.000
Positive	19	44			
Negative	31	6			


*Predictive Factors Study*

The results obtained (Figure 1 and Table 5) show that serum MDA, Erythrocytic GSH, and Leukocytic SOD levels are the highest percentage of specificity (100%) and important

percentage of sensitivity (50.0, 43.8, 75%) respectively for women with osteoporosis. In addition, Serum TAC, Leukocyte GSH, and Leukocytic catalase were significant predictive factors for osteoporosis in postmenopausal women.

**Table 5.** Sensitivity, Specificity and AUC Values of Biological Markers for Women with Osteoporosis

Test Result Variable(s)	Sensitivity (%)	Specificity (%)	AUC (%)	SE	CI <sub>95%</sub>	P-value
Serum TAC	50.0	25	19.1	0.076	0.043 0.340	<b>0.003</b>
Leukocytic SOD	75.0	100	62.5	0.063	0.001 0.249	<b>0.000</b>
Serum SOD	68.8	31.2	32.2	0.099	0.129 0.516	0.086
Erythrocytic GSH	43.8	100	100	0.000	0.000 0.000	<b>0.000</b>
Leukocyte GSH	18.8	25	10.5	0.058	0.000 0.219	<b>0.000</b>
Serum GSH	81.3	12.5	42.8	0.103	0.226 0.630	0.486
Serum vitamin C	43.8	62.5	58.6	0.103	0.384 0.788	0.407
Leukocytic catalase	50.0	62	15.8	0.071	0.019 0.298	<b>0.001</b>
Serum catalase	31.3	43.7	30.9	0.096	0.121 0.496	0.065
Erythrocytic MDA	43.8	25	34.0	0.100	0.145 0.535	0.122
Leukocytic MDA	43.8	100	70.7	0.098	0.515 0.899	0.064
Serum MDA	50.0	100	99.6	0.006	0.983 1.000	<b>0.000</b>



**Figure 1.** ROC Curve for Biological Markers in Women with Osteoporosis

*Correlation between Biological Markers*

The results (Table 6) represent the correlation between oxidative stress parameters (serum ORAC, leukocyte SOD, erythrocyte GSH, leukocyte GSH, leukocyte catalase, and serum

MDA), hematological parameters (WBC, HGB, RBC, HCT, PLT), and biochemical parameters (calcium, ALP) in patients group with osteoporosis. There was a positive correlation (P<0.05) between erythrocyte GSH and calcium (P=0,005 and R=0,381),

erythrocyte GSH and ALP ( $P=0,047$  and  $R=0,393$ ), Leukocyte GSH and serum calcium ( $P=0,046$  and  $R=0,34$ ). There was no

correlation ( $P>0.05$ ) between the rest of the correlation test in the patients' groups.

**Table 6.** Correlation between Biological Markers for Women with Osteoporosis

Parameters		WBC	HGB	RBC	HCT	PLT	Serum Calcium	Serum ALP
Serum ORAC	P	0,926	0.565	0,729	0,642	0,429	0,447	0,734
	R	0,019	0.118	0,071	0,096	0,162	-0,108	0,070
Leukocyte SOD	P	0,580	0,879	0,582	0,420	0,644	0,435	0,146
	R	-0,114	-0,031	-0,113	-0,165	-0,095	0,111	-0,293
Erythrocyte GSH	P	0,624	0,221	0,665	0,303	0,617	<b>0,005</b>	<b>0,047</b>
	R	-0,101	-0,248	-0,089	-0,210	0,103	0,381	0,393
Leukocyte GSH	P	0,458	0,463	0,929	0,700	0,320	<b>0,046</b>	0,955
	R	0,152	-0,151	-0,018	-0,079	0,203	0,340	-0,012
Leukocyte Catalase	P	0,318	0,379	0,962	0,722	0,761	0,375	0,115
	R	-0,204	-0,180	-0,010	-0,073	-0,063	-0,126	0,317
Serum MDA	P	0,938	0,870	0,715	0,619	0,285	0,582	0,521
	R	0,016	0,034	-0,075	0,102	-0,218	0,078	-0,132

The obtained results show a significant increase in PAL level in the patient group as compared to control. This result is in line with the research of Pardhe *et al.*, (2017), who show that the PAL level was slightly higher in the post-menopausal group. All Bone Mineral Density (BMD) results were remarkably reduced by PAL increment, while bone-specific alkaline phosphatase, is an indicator of bone turnover and formation and is utilized in the skeletal status evaluation (Hailing *et al.*, 2018). On the other hand, we found a significant decrease in calcium level in the patients' group as compared to the control group. This result is consistent with the research of (Beto, 2015), whose findings demonstrated that the serum calcium level was significantly lower in the post-menopausal group. Calcium plays a key role in human physiology. As a main constituent of the mineral component calcium provides stiffness to the collagen network of the mature bone. Inadequate calcium accrual, resulting in a sub-optimal bone mass peak and low bone mineralization, is a significant parameter favoring osteoporosis and fracture (Kelly *et al.*, 2020). The results of hematological parameters indicated a remarkable increment in WBC, HGB, HCT, and PLT in the osteoporosis patients group in comparison to the control group. These results support a possible linkage between bone metabolism and hematopoiesis. Hematopoiesis is the process by which immature blood cells develop into mature cells (Paspaliaris and Kolios, 2019). According to Schyrr *et al.*, (2018), differences in the osteoporotic bone microenvironment translate into altered dynamics upon hematopoietic stress. Moreover, Valderrábano *et al.*, (2018) found that low bone health would result in enhanced cells of myeloid lineage such as neutrophils and monocytes. Osteoblastic lineage cells might affect neutrophils and monocytes differentiation. The enhancement in neutrophils can be associated with the chronic inflammation that happens with aging. The results of the oxidative stress study showed for the osteoporosis patients group a very high significant increase in serum MDA level as compared to the control women. The results showed that serum MDA has a high specificity in ROC statistics, which showed the importance of MDA in the prognostic of osteoporosis. The results found were similar to those observed in the study of Berköz *et al.* (2017) which showed that serum MDA levels were significantly higher

in postmenopausal women with osteoporosis than in the healthy controls. Sakuraba *et al.*, (2020) reported that MDA had an osteoclastic activity. Our results show that leukocyte MDA is significantly increased in the osteoporosis patients group as compared to control women, with high specificity in the ROC statistic test, which showed the importance of this parameter in the identification of the disease. These results are supported by Ahmedian *et al.*, (2017) study. Raghavan *et al.* (2012) found that MDA could significantly induce key inflammatory cytokine expression in lymphocyte via oxidant stress, signaling pathways (p38MAPK), and transcriptional factors (NF- $\kappa$ B), which in turn enhance lymphocyte activation. The results also show that for the osteoporosis patients group, a very high significant decrease in leukocyte and serum SOD level ( $P<0.001$ ) in comparison to that in the control. Depressing actions of the antioxidant enzymes such as SOD indicated a defense mechanism that has been overwhelmed in mitigating the enhanced superoxide production by the osteoclasts showed by enhanced contents of MDA in the serum (Kuyumcu and Aycan, 2018) and it might cause markedly increased bone demineralization and, as a result, may increase destructive free radical levels (Bacou *et al.*, 2021). The results show that for the osteoporosis patients group, a highly significant decrease in serum TAC level ( $P<0.01$ ) as compared to that in the control. TAC method is relevant to in vivo conditions since it uses a biologically relevant free radical source (peroxyl radical) that is the highest common free radical in human biology. It considers both degrees of inhibition and inhibition time of free radical action as a result of antioxidants (Hunyadi, 2019). The results of the oxidative stress study showed that there is a significant decrease in GSH, catalase, and SOD level of WBC, in GSH level of RBC, and both of GSH and catalase level of serum in the patients' group compared to controls. Glutathione (GSH) is a non-enzymatic antioxidant that aids the defense mechanism against oxidative stress created by free radicals (Derouiche *et al.*, 2019). To reduce the cell-damaging impacts of ROS (reactive oxygen species), aerobic organisms evolved by expressing different antioxidant defenses, such as catalase. The mechanisms by which cells sense  $H_2O_2$  and  $O_2^\bullet$  are not comprehended; however, many transcriptional parameters that adjust the

expression of antioxidant genes are adjusted by decrease counteractions and oxidation (Tonelli *et al.*, 2018). Intracellular Redox Imbalance resulting from SOD shortage has a vital impact on the progress and development ion of bone fragility both in vitro and in vivo (Tan *et al.*, 2018).

### CONCLUSION

This study indicates that hematological parameters change and Oxidative stress were correlated with the osteoporosis disease of menopause in women as a reason or as a developmental factor for it. And through the ROC analysis results MDA, GSH, SOD, and TAC were considered to be the most important markers which contribute in the early detection of osteoporosis disease in post-menopause women.

**ACKNOWLEDGMENTS:** This research was made possible by the D01N01UN390120190001 research project, which was funded by the Algerian Ministry of Higher Education and the Directorate General for Scientific Research and Technological Development.

**CONFLICT OF INTEREST:** None

**FINANCIAL SUPPORT:** None

**ETHICS STATEMENT:** Ethical approval was obtained from the ethics committee (28 EC/DCMB/FNSL/EU2020) of the department of cellular and molecular biology, Faculty of Natural Sciences and Life, University of El Oued.

### REFERENCES

- Aebi, H. (1984). Catalase in vitro. *Methods in Enzymology*, 105(1), 121-126. doi: 10.1016/s0076-6879(84)05016-3.
- Ahmadian, F., Mozaffari-Khosravi, H., Azarain, M. H., Faraji, R., & Zavar-Reza, J. (2017). The effect of consumption of garlic tablet on proteins oxidation biomarkers in postmenopausal osteoporotic women: A randomized clinical trial. *Electronic Physician*, 9(11), 5670-5675.
- Al-Shali, K. Z., Alama, M. N., Ajabnoor, M. A., Al-Fayaz, F. F., & Balubaid, K. H. (2019). Osteoprotegerin Levels among Saudi Cardiovascular Patients. *International Journal of Pharmaceutical Research & Allied Sciences*, 8(1), 95-101.
- Atoussi, O., Chetehouna, S., Boulaares, I., Guemari, I. Y., & Derouiche, S. (2021). Analysis of Blood Pressure, Lipid Profile and Hematological Biomarkers in men Addicted to Tobacco Chewing. *Research Journal of Pharmacology and Pharmacodynamics*, 13(1), 1-5.
- Bacou, E., Walk, C., Rider, S., Litta, G., & Perez-Calvo, E. (2021). Dietary Oxidative Distress: A Review of Nutritional Challenges as Models for Poultry, Swine, and Fish. *Antioxidants*, 10(4), 525.
- Beauchamp, C., & Fridovich I. (1971). Superoxide dismutase: Improved assays and an assay applicable to acrylamide gels. *Analytical Biochemistry*, 44(1), 276-287.
- Berköz, M., Sağır, Ö., Yalın, S., Çömelekoğlu, Ü., Sögüt, F., & Eroğlu, P. (2017). Effect of postmenopausal strontium ranelate treatment on oxidative stress in rat skin tissue. *Eastern Journal of Medicine*, 22(4), 162-166.
- Beto, J. A. (2015). The role of calcium in human aging. *Clinical Nutrition Research*, 4(1), 1-8.
- Chetehouna, S., Atoussi, O., Boulaares, I., Guemari, I. Y., & Derouiche, S. (2020). The effect of Chronic Tobacco smoking on Atherogenic index and Cardiovascular diseases risk in El-Oued (Algeria) men. *Asian Journal Research in Chemistry*, 13(6), 1-5.
- Derouiche, S., Abbas, K., & Djermoune, M. (2017). Polysaccharides and ascorbic acid content and the effect of aqueous extract of Portulaca oleracea in high-fat diet-induced obesity, dyslipidemia and liver damage in albino wistar rats. *Algerian Journal Arid Environnement*, 7(2), 16-26.
- Derouiche, S., Atoussi, N., & Guediri, S. (2019). Assessment of Hematological Parameters, Enzymes Activities, and Oxidative Stress Markers in Salivary and Blood of Algerian Breast Cancer Patients Receiving Chemotherapy. *Journal of Biochemistry and Technology*, 10(4), 50-58.
- Hailing, C., Jufen, L., & Qian, W. (2018). Associations between bone-alkaline phosphatase and bone mineral density in adults with and without diabetes. *Journal List Medicine (Baltimore)*, 97(17), 1-7. doi: 10.1097/MD.00000000000010432
- Hunyadi A. (2019). The mechanism(s) of action of antioxidants: From scavenging reactive oxygen/nitrogen species to redox signaling and the generation of bioactive secondary metabolites. *Medicinal Research Reviews*, 39(6), 2505-2533.
- Jagota, S. K., & Dani, H. M. (1982). New colorimetric technique for the estimation of vitamin C using Folin phenol reagent. *Analytical Biochemistry*, 127(1), 178-182.
- Kelly, R. R., Sidles, S. J., & LaRue, A. C. (2020). Effects of Neurological Disorders on Bone Health. *Frontiers in Psychology*, 11, 612366.
- Kuyumcu, F., & Aycan, A. (2018). Evaluation of Oxidative Stress Levels and Antioxidant Enzyme Activities in Burst Fractures. *Medical Science Monitor: International Medical Journal of Experimental and Clinical Research*, 24, 225-234. doi: 10.12659/msm.908312
- Nambi, G., Abdelbasset, W. K., Moawd, S. A., Sakr, H. R., Elnegamy, T. E., & George, J. S. (2020). Effect of virtual reality training on post-menopausal osteoporotic women. *Archives of Pharmacy Practice*, 11(2), 6-12.
- Oyaizu, M. (1986). Studies on products of browning reaction: antioxidant activities of products of browning reaction prepared from glucosamine. *Japanese Journal of Nutrition*, 44(6), 307-315.
- Pardhe, B. D., Pathak, S., Bhetwal, A., Ghimire, S., Shakya, S., Khanal, P. R., & Marahatta, S. B. (2017). Effect of age and estrogen on biochemical markers of bone turnover in postmenopausal women: a population-based study from Nepal. *International Journal of Women's Health*, 9, 781-788.
- Paspaliaris, V., & Kolios G. (2019). Stem cells in Osteoporosis: From Biology to New Therapeutic Approaches. *Stem Cells International*, 2019. doi: 10.1155/2019/1730978
- Raghavan, S., Subramaniam, G., & Shanmugam, N. (2012). Proinflammatory effects of malondialdehyde in lymphocytes. *Leukocyte Biology*, 92(5), 1055-1067.
- Sakuraba, K., Krishnamurthy, A., Circiumaru, A., Sun, M., Joshua, V., Engström, M., Zheng, X., Xu, C., Amara, K., Malmström, V. et al. (2020). SAT0017 Metabolic changes induced by anti-malondialdehyde/malindialdehyde-acetaldehyde antibodies promote osteoclast development. *Annals of the Rheumatic Diseases*, 79, 938-939.

- Sastre, J., Pallardo, F. V., Corcia dela Asuncion, J., & Vina, J. (2000). Mitochondria, oxidative stress and aging. *Free Radical Research*, 32(3), 189-198. doi: 10.1080/10715760003002016/50009-6
- Schyrer, F., Wolfer, A., Pasquier, J., Nicoulaz, A.L., Lamy, O., & Naveiras, O. (2018). Correlation study between osteoporosis and hematopoiesis in the context of adjuvant chemotherapy for breast cancer. *Annals of Hematology*, 97(1), 309-317.
- Sözen, T., Özışık, L., & Başaran, N. Ç. (2017). An overview and management of osteoporosis. *European Journal of Rheumatology*, 4(1), 46-56.
- Tan, B. L., Norhaizan, M. E., Liew, W. P., & Sulaiman Rahman, H. (2018). Antioxidant and Oxidative Stress: A Mutual Interplay in Age-Related Diseases. *Frontiers in Pharmacology*, 9, 1162.
- Tomlinson, R. E., Christiansen, B. A., Giannone, A. A., & Genetos D.C. (2020). The Role of Nerves in Skeletal Development, Adaptation, and Aging. *Frontiers in Endocrinology*, 11, 646
- Tonelli, C., Chio, C., & Tuveson, D. A. (2018). Transcriptional Regulation by Nrf2. *Antioxidants & Redox Signaling*, 29(17), 1727-1745.
- Valderrábano, R. J., Lui, L. Y., Lee, J., Cummings, S. R., Orwoll, E. S., Hoffman, A. R., & Wu, J. Y. (2018). Bone Density Loss Is Associated with Blood Cell Counts. *Bone and Mineral Research*, 32(2), 212-220. doi: 10.1002/jbmr.3000
- Weak, B. G., & Cory, J. G. (1988). Ribonucleotide reductase activity and growth of glutathione-depleted mouse L1210 cells in vitro. *Cancer Letters*, 40(3), 257-264.
- Weiss, G., Skurnick, J. H., Goldsmith, L. T., Santoro, N. F., & Park, S. J. (2004). Menopause and hypothalamic-pituitary sensitivity to estrogen. *The Journal of the American Medical Association*, 292(24), 2991-2996. doi:10.1001/jama.292.24.2991.
- Wright, N. C., Looker, A. C., Saag, K. G., Curtis, J. R., Delzell, E. S., Randall, S., & Dawson-Hughes, B. (2014). The recent prevalence of osteoporosis and low bone mass in the United States based on bone mineral density at the femoral neck or lumbar spine. *Journal of Bone and Mineral Research*, 29(11), 2520-6.
- Zeidabadi, A., Yazdanpanahi, Z., Dabbaghmanesh, M. H., Sasani, M. R., Emamghoreishi, M., & Akbarzadeh, M. (2020). The effect of Salvia officinalis extract on symptoms of flushing, night sweat, sleep disorders, and score of forgetfulness in postmenopausal women. *Journal of Family Medicine and Primary Care*, 9(2), 1086-1092.
- Zerzour, A., Heddig, N. E., & Samir Derouiche, S. (2020). Analysis of Osteoporosis risk factors in Menopausal women's of Algeria population. *Asian Journal of Research in Pharmaceutical Sciences*, 10(2), 79-84.