



## Isolation and Optimization Growth Parameters for Enhanced Bioactivity Profiles of Streptomyces Strain Isolated from Saudi Arabia

Afra Mohammed Baghdadi<sup>1\*</sup>

<sup>1</sup>Department of Biology, College of Sciences, University of Jeddah, Jeddah, Saudi Arabia.

### ABSTRACT

The genus *Streptomyces* is a unique subgroup of actinomycetes bacteria that is well-known as prolific producers of many bioactive secondary metabolites as well as antibiotics. *Streptomyces globosus* (*S. globosus*) is a bacterium species from the genus of *Streptomyces*. Objective: This study aimed to optimize the growth conditions required for the maximum antibacterial activity of *S. globosus* isolated from soil samples in Saudi Arabia. The effect of dark/light, temperature, and pH on the growth and antibacterial effect against *E. coli* ATCC35218, *E. coli* ATCC25922, *Salmonella* sp., *P. aeruginosa* ATCC27853, *S. aureus* ATCCBAA977, and *St. pneumonia* ATCC49619 were optimized. The results of this study showed that the highest antibacterial activity was at 30 °C after 7 days of incubation.

Furthermore, the best to produce antibacterial activity was under alkaline conditions (pH 8.5). Besides, the isolate didn't show antibacterial activity at the shaking incubator. This study showed that all organic solvents (ethyl-alcohol, methanol, chloroform, ethyl-acetate, and acetone) successfully extracted the antibacterial compound. Still, the compounds extracted by chloroform had the highest antibacterial activity. The MIC of the chloroform extract was against *S. aureus* ATCCBAA977. The study described the optimized condition for best *S. globosus* activity. The *S. globosus* showed a broad spectrum of antimicrobial activities against the test organisms. This opened further research investigations on the purification and structural characterization of the active compounds from the crude extract.

**Keywords:** *Streptomyces globosus*, Condition, Dark/light, Antibacterial temperature activity, pH

**Corresponding author:** Afra Mohammed Baghdadi

**e-mail** ✉ [amboghdadi@uj.edu.sa](mailto:amboghdadi@uj.edu.sa)

**Received:** 08 December 2021

**Accepted:** 04 May 2022

### INTRODUCTION

Antimicrobial is used to spare people from contamination. It is a substance or a synthetic item with an inhibitory or destructive impact on the microorganisms (Pidcock, 2012; Aloqayli *et al.*, 2022). The advancement of safe strains is a characteristic wonder that happens through choice weight on the antimicrobial microorganism populace (Chellat *et al.*, 2016). Nonetheless, some of these antimicrobials have been rendered dormant by resistant systems (Nikolaidis *et al.*, 2014; Batarseh *et al.*, 2022). Several natural antimicrobials were extracted from soil microbes (Jain & Pundir, 2011; Qinn *et al.*, 2020).

Actinomycetes are a group of prokaryotic microorganisms, gram-positive bacteria with high guanine-cytosine in their DNA (Adegboye & Babalola, 2012). They are considered biotechnologically important organisms since they are responsible for producing about half of the bioactive secondary metabolites, including antibiotics. They are filamentous bacteria that produce two types of branching mycelium, namely aerial and substrate mycelium (Yuan *et al.*, 2010).

*Streptomyces* (family *Streptomycetaceae*) are the most significant Actinobacteria (Rong *et al.*, 2009; Radwan *et al.*, 2022). It is disseminated in soil, water, and other indigenous habitats (Singh *et al.*, 2006). In all years, *Streptomyces* strains as a wellspring of various bioactive mixes, for example, anthelmintic chemicals, herbicides, hostile to malignancy

drugs, development components, and insusceptible modulators (Kumar *et al.*, 2020; Adeyemo *et al.*, 2021).

Factors influencing the number and types of actinomycetes present in a particular soil are geographical location, such as soil type, temperature, organic matter content, moisture content, cultivation, and aeration. Actinomycetes act as a major component of the microbial population in most soil. About 90% of the total actinomycetes population consists of *Streptomyces* species (Harir *et al.*, 2018). Almost 80% of the world's antibiotics are known to come from actinomycetes, mostly from the genera *Streptomyces* and *Micromonospora* (Pandey *et al.*, 2004).

A major global healthcare problem has occurred in the 21<sup>st</sup> century because of the increasing resistance of microorganisms against commonly used antibiotics. A renewed search for new antimicrobial agents from *Streptomyces* has occurred because of this rapid emergence of antimicrobial resistance among pathogenic microorganisms (Alanis, 2005; Sebak *et al.* 2021).

Microscopic organisms are profoundly versatile pathogens that create protection from antimicrobials through a few mechanisms. The creation of  $\beta$ -lactamases that hydrolyze the  $\beta$ -lactam ring is the most widely recognized resistant system for these microscopic organisms against  $\beta$ -lactam antimicrobials (Pfeifer *et al.*, 2010; Tooke *et al.*, 2019). Creating broad range  $\beta$ -lactamases (ESBLs) is a noteworthy opposition instrument that obstructs the antimicrobial treatment of contaminations brought about by Enterobacteriaceae and is a genuine risk to the presently accessible antimicrobial arsenal (Shaikh *et al.*, 2015). Bacterial efflux pumps effectively transport numerous

antimicrobials out of the cell. They are significant contributors to the characteristic resistance of gram-negative microscopic organisms to many medications that can be utilized to treat gram-positive bacterial contaminations (Blair *et al.*, 2015). This study aimed to optimize the growth conditions required for the maximum antibacterial efficacy of *Streptomyces globosus* (*S. globosus*) separated from soil samples in Saudi Arabia.

## MATERIALS AND METHODS

### Effect of different pH values

The 100 ml Calibration of sterile starch-nitrate broth medium for different pH levels; 6, 6.5, 7, 7.5, 8, 8.5, 9, and 9.5 using Na OH for alkaline and HCl for acidic before sterilizing medium was performed. The cultures were incubated at 30°C. The culture filtrates were centrifuged (4000 rpm in 30 m). The antibacterial efficacy of CFF was investigated using the agar well diffusion technique. Twenty-four hours post-incubation (35 ± 2°C), the antibacterial efficacy was calculated by observing the zone of inhibition radius in millimeters.

### Effects of the light factor

Nine Petri plates by the selected *streptomyces* isolate were incubated (three in light, dark, or 12 h light/ 12 h dark) at 28°C for 7 days. Two agar discs (6 diameters) of *Streptomyces* isolate were placed over the surface of the tested bacteria's cultures with equal distance and incubated at 35 ± 2 °C for one day. After incubation, the antibacterial activity was estimated by measuring the inhibition zones' diameter in (mm).

### Effect of different incubation periods at different incubation temperatures

The chosen *Streptomyces sp.* was cultured on sterile starch-nitrate broth (100 ml) medium at different temperatures (25°, 30°, 35° and 40° C) for various periods (1, 2, 3, 4, 5, 6, 7, 8 and 9). At each incubation period and different temperatures, the culture filtrates were centrifuged (4000 rpm in 30 m). The antibacterial efficacy of cell-free filtrate (CFF) was investigated using the agar well diffusion technique. Twenty-four hours post-incubation (35 ± 2 °C), the antibacterial efficacy was calculated by observing the zone of inhibition radius in millimeters.

### Effect of shaking incubator

The chosen *Streptomyces sp.* was cultured on sterile starch-nitrate broth (100 ml) medium in optimum conditions. Half of the cultures were incubated on a shaking incubator, and others were incubated in a constant incubator. The culture filtrates

were centrifuged (4000 rpm in 30 m). The antibacterial efficacy of CFF was investigated using the agar well diffusion technique. Twenty-four hours post-incubation (35 ± 2 °C), the antibacterial efficacy was calculated by observing the zone of inhibition radius in millimeters.

### Extraction of an antibacterial compound with organic solvents

Four different solvents (ethyl-alcohol, methanol, chloroform, ethyl-acetate, and acetone) were used to extract the antibacterial activity agent that was produced by the selected *Streptomyces sp.* The chosen *Streptomyces sp.* was cultured on sterile starch-nitrate broth (100 ml) medium in optimum conditions. The culture filtrates were centrifuged (4000 rpm in 30 m). The CFF was transferred aseptically into the conical flasks, and an equal volume of solvents was separately added to the CFF and shaken for 20 min. After gathering the organic phase, the solvent was evaporated under a vacuum. 50 µl of the extract was used to investigate antibacterial efficacy using the agar well diffusion technique. Twenty-four hours post-incubation (37 °C), the antibacterial efficacy was calculated by observing the zone of inhibition radius in millimeters.

### MIC values against nosocomial infection bacteria

The process includes the preparation of a double dilution of the antimicrobial agents in a liquid medium. Upon diluting the standardized bacterial culture, each tube with a bacterial suspension was packed in the same medium. The inoculated tubes were incubated (mostly without agitation) under suitable conditions depending upon the test microorganism.

### Statistics

The data presented as a mean of 3 replicates ± SD. The difference between the means was statistically compared using Microsoft Excel software.

## RESULTS AND DISCUSSION

### Effect of the light factor on antibacterial activity of *S. globosus*

**Table 1** revealed no significant difference in the antibacterial activity when *S. globosus* was grown in different lighting. It had antibacterial activity against all the test bacteria except *P. earuginosa* ATCC27853, but no differences in the antibacterial activity against the other test bacterial isolates when *S. globosus* was grown in light, dark, or 12h light / 12h dark, it showed antibacterial activity against *E.coli* ATCC35218 only when it was grown in dark.

**Table 1.** Effect of the light factor on antibacterial activity of *S. globosus*

Light factors	Inhibition zone diameter (mm)					
	<i>E. coli</i> ATCC35218	<i>E. coli</i> ATCC25922	<i>Salmonella</i> <i>sp.</i>	<i>P. earuginosa</i> ATCC27853	<i>S. aureus</i> ATCCBAA977	<i>St. pneumonia</i> ATCC49619
Dark	11.3 ± 0.47	15.6 ± 0.47	18.3 ± 0.47	-	24.3 ± 0.47	14.6 ± 0.47
Light	-	15.3 ± 0.47	17.16 ± 0.23	-	23.16 ± 0.23	13.3 ± 0.23
12h light / 12h dark	-	15.3 ± 0.47	17.83 ± 0.23	-	24 ± 0.00	14.3 ± 0.47


### Effect of different incubation periods and different incubation temperatures on antibacterial activity of *S. globosus*

Incubation of *S. globosus* at 25°C showed antibacterial activity against *S. aureus* ATCCBAA977 only after five days of incubation and increased by increasing the incubation period. The most

excellent antibacterial activity was after seven days of incubation, and the inhibition zone was 14.3 mm. Still, there was no antibacterial activity against other Gram-positive and Gram-negative bacteria (**Figure 1**).

Results in **Figures 1a-1c** revealed that antibacterial activity against most test bacteria was noticed and increased with

increasing the incubation period at 30°C. The optimum antibacterial activity was after seven days of incubation. The diameters of inhibition zones were (13.3 mm, 18.0 mm, 12.0 mm, 19.3 mm, and 16.3 mm) against *E. coli* ATCC35218, *E. coli* ATCC25922, *Salmonella* sp., *S. aureus* ATCCBAA977 and *St. pneumonia* ATCC49619, respectively.



**Figure 1.** Effect of different incubation periods and different incubation temperatures on antibacterial activity of *S. globosus*

From the results in **Figure 1**, when *S. globosus* was incubated at 35°C, the antibacterial efficacy was versus *E. coli* ATCC25922, *Salmonella* sp., and *S. aureus* ATCCBAA977. The antibacterial activity against *Salmonella* sp. and *S. aureus* ATCCBAA977 began from the third day but after five days produced antibacterial


activity against *E. coli* ATCC25922. The optimum antibacterial activity was on the fifth day, and the diameters of inhibition zones were (12.6 mm, 15.0 mm, and 16.3 mm) versus *E. coli* ATCC25922, *Salmonella* sp, and *S. aureus* ATCCBAA977. *S.*

*globosus* at 40°C did not show antibacterial activity against all test bacteria.

*Impact of various pH conditions on antibacterial efficacy of S. globosus*

*S. globosus* produce antibacterial activity against *S. aureus* ATCCBAA977 at pH 6, against *E. coli* ATCC25922, and *S. aureus* ATCCBAA977 at pH 6.5, and against all test bacterial isolate


except *P. earuginosa* ATCC27853 at pH 7. **Figure 2** shows that the antibacterial activity of *S. globosus* against all test bacteria increased by increasing the fermentation medium's pH values. The highest antibacterial activity was at pH 8.5, which was decreased above. It was noticed that *S. globosus* produced antibacterial activity against all test isolates and the highest was against *S. aureus* ATCCBAA977. The inhibition zone diameter was (20.3 mm) and decreased slowly above.



**Figure 2.** Impact of various pH efficacy of *S. globosus*. condition on antibacterial

*Effect of shaking incubator on antibacterial activity of S. globosus*  
Results in **Figure 3** show the difference between the antibacterial activity produced by *S. globosus* when the cultures of this isolate were incubated in a constant and shaking

incubator. When incubated in a shaking incubator, *S. globosus* can't have antibacterial activity against all the test bacteria. Different antibacterial activities were produced when cultures of *S. globosus* were incubated in a constant incubator.




**Figure 3.** Effect of shaking incubator on antibacterial activity of *S. globosus*.

The diameters of inhibition zones were (15.3 mm, 14.0 mm, 13.3 mm, 14.3 mm, 24.0 mm, and 15.3 mm) against *E. coli* ATCC35218, *E. coli* ATCC25922, *Salmonella* sp., *P. earuginosa* ATCC27853, *S. aureus* ATCCBAA977 and *St. pneumonia* ATCC49619, respectively.

**Figure 4** revealed that all organic solvents succeeded in extracting the antibacterial compound. Better extraction was obtained using chloroform. The diameters of inhibition zones were (17.6 mm, 16.3 mm, 17.3 mm, 15.6 mm, 27.6 mm, and 17.3 mm) against *E. coli* ATCC35218, *E. coli* ATCC25922, *Salmonella* sp., *P. earuginosa* ATCC27853, *S. aureus* ATCCBAA977 and *St. pneumonia* ATCC49619, respectively.

*Effect of different organic solvents used in the extraction of the antibacterial compound from S. globosus on its antibacterial activity.*



**Figure 4.** Effect of different organic solvents used to extract the antibacterial compound from *S. globosus* on its antibacterial activity.

MIC values of the chloroform extract against nosocomial infection bacteria

The MIC values against all the tested bacteria showed different results. *S. aureus* ATCCBAA977 has been inhibited at (16 µg/ml), which was the lowest concentration for inhibition. *E. coli*

ATCC35218, *Salmonella* sp., and *St. Pneumonia* ATCC49619 bacteria have been inhibited at (32 µg/ml). *E. coli* ATCC25922 inhibited at (64 µg/ml). The highest concentration for inhibition was at (125 µg/ml) against *P. earuginosa* ATCC27853 (**Table 2**).

**Table 2.** MIC values of the chloroform extract against nosocomial infection bacteria.

Nosocomial infection bacteria	Minimum inhibitory concentration (MIC)					
	<i>E. coli</i> ATCC35218	<i>E. coli</i> ATCC25922	<i>Salmonella</i> sp.	<i>P. earuginosa</i> ATCC27853	<i>S. aureus</i> ATCCBAA977	<i>St. pneumonia</i> ATCC49619
Chloroform	32	64	32	125	16	32

In this work, *Streptomyces* sp. was dominantly separated from the soil specimen on a starch-nitrate medium. This finding is congruent with earlier studies, which reported that the recovery percentage of *Streptomyces* sp. with antimicrobial activity was higher than other actinobacterial genera (Kitouni *et al.*, 2005). Rare actinobacteria are also important in discovering novel antibiotics (Tiwari & Gupta, 2012).

This study aimed to optimize the culture condition, which will yield the most optimum antimicrobial efficacy of *Streptomyces* sp. The incubation temperature can influence bacterial growth and the development of biosynthetic pathways of the metabolites (Stanbury, 2016). Also, Pandey *et al.* (2005) and Vijayakumar *et al.* (2012) showed that for the optimum production of antibiotics, certain carbon and nitrogen sources are required study done by Vasavada *et al.* (2006) showed that the use of different media, pH, salinity, and carbon and nitrogen affect the growth and antibiotic production by Actinomycetes. Results of the impact of incubation time and temperature on the optimum efficacy of *S. globosus* showed that the highest antibacterial activity was at 30 °C after 7 days of incubation. This result is similar to Sujatha *et al.* (2005), with the highest antimicrobial activity at 30 °C. On the other hand, Ripa *et al.* (2009) reported that isolated *Streptomyces* sp. produced high antibiotic levels when incubated at 39 °C.

This study showed that the best to produce antibacterial activity was under alkaline conditions (pH 8.5). Similar results had been reported earlier by Basilio *et al.* (2003). Ripa *et al.* (2009) showed that the pH 8.0 was optimum for antibiotics produced by *Streptomyces* sp. This may be attributed to the presence of active enzymes for antimicrobial metabolites synthesis at pH 7-

9. Each *Streptomyces* strain has an optimum, minimum, and maximum pH at which it gave an optimum enzyme activity (Vijayakumar *et al.*, 2012).

This study showed that the isolate didn't show antibacterial activity at the shaking incubator. Gottschalk *et al.* (2003), in their research using *S. viridosporus*, suggested that the rise in the rate of agitation up to 500 rpm contributed to a persistent decline in the growth and development of cells. A rise in the stirring rate may have contributed to high shearing outcomes that were harmful to the solidity of the fungi.

This study showed that all organic solvents (ethyl-alcohol, methanol, chloroform, ethyl-acetate, and acetone) successfully extracted the antibacterial compound. Still, the compounds extracted by chloroform had the highest antibacterial activity. Our results were similar to those of Sathi *et al.* (2001). Disagree with our results, Vijayakumar *et al.* (2012) confirmed that extraction of *Streptomyces* sp. with ethyl acetate produced more activity by generating the maximum inhibition versus *P. Vulgaris* and minimum inhibition versus *S. aureus*. Besides, Balachandran *et al.* (2012) showed that extraction of *Methylobacterium* sp. With ethyl acetate produced the maximum inhibition versus *K. pneumonia* and minimum inhibition versus *B. subtilis*.

## CONCLUSION

These findings indicated that the soil samples of the Al-Baha region have actinomycetes with metabolites that control bacterial pathogens. The extracted metabolites obtained from

the isolate of our study showed very good antimicrobial activity against four different clinical pathogens. The further structural characterization of these molecules promotes the utilization of this active principle as a lead in case of drug discovery from soil-based formulations to eradicate the emerging infectious drug-resistant pathogenic microorganisms.

**ACKNOWLEDGMENTS:** None

**CONFLICT OF INTEREST:** None

**FINANCIAL SUPPORT:** None

**ETHICS STATEMENT:** None

## REFERENCES

- Adegboye, M. F., & Babalola, O. O. (2012). Taxonomy and ecology of antibiotic producing actinomycetes. *African Journal of Agricultural Research*, 7(15), 2255-2261.
- Adeyemo, O. M., Ja'afaru, M. I., & Adams, F. V. (2021). Isolation, characterization, antimicrobial and other bioactivity profiles of three Streptomyces strains isolated from Lake Gerio, Yola, Adamawa State, Nigeria. *Bulletin of the National Research Centre*, 45(1), 1-13. doi:10.1186/s42269-021-00606-x
- Alanis, A. J. (2005). Resistance to antibiotics: Are we in the post-antibiotic era? *Archives of Medical Research*, 36(6), 697-705.
- Aloqayli, M. A., Alshuwirikh, R. M., Alghwery, N. S., Ansari, S. H., & Alhussain, B. S. (2022). Quality assessment of composite restorations performed by the dental students: A retrospective study. *Annals of Dental Specialty*, 10(2), 101-105.
- Balachandran, C., Duraipandiyar, V., Al-Dhabi, N. A., Balakrishna, K., Kalia, N. P., Rajput, V. S., Khan, I. A., & Ignacimuthu, S. (2012). Antimicrobial and antimycobacterial activities of methyl caffeate isolated from *Solanum torvum* Swartz. fruit. *Indian Journal of Microbiology*, 52(4), 676-681.
- Basilio, A., Gonzalez, I., Vicente, M. F., Gorrochategui, J., Cabello, A., Gonzalez, A., & Genilloud, O. (2003). Patterns of antimicrobial activities from soil actinomycetes isolated under different conditions of pH and salinity. *Journal of Applied Microbiology*, 95(4), 814-823.
- Batarseh, N., Khalil, R., & Al-Domi, H. A. (2022). Hypothalamic neuroinflammation induced by obesity and the effect of Liraglutide. *Journal of Advanced Pharmacy Education and Research*, 12(1), 46-55.
- Blair, J., Webber, M. A., Baylay, A. J., Ogbolu, D. O., & Piddock, L. J. (2015). Molecular mechanisms of antibiotic resistance. *Nature Reviews Microbiology*, 13(1), 42-51.
- Chellat, M. F., Raguž, L., & Riedl, R. (2016). Targeting antibiotic resistance. *Angewandte Chemie International Edition*, 55(23), 6600-6626.
- Gottschalk, L. M., Nobrega, R., & Bon, E. P. (2003). Effect of aeration on lignin peroxidase production by *Streptomyces viridosporus* T7A. In *Biotechnology for Fuels and Chemicals* (pp. 799-807). Humana Press, Totowa, NJ.
- Harir, M., Bendif, H., Bellahcene, M., Fortas, Z., & Pogni, R. (2018). Streptomyces secondary metabolites. *Basic Biology and Applications of Actinobacteria*, 6, 99-122.
- Jain, P., & Pundir, R. K. (2011). Effect of fermentation medium, pH and temperature variations on antibacterial soil fungal metabolite production. *Journal of Agricultural Technology*, 7(2), 247-269.
- Kitouni, M., Boudemagh, A., Oulmi, L., Reghioia, S., Boughachiche, F., Zerizer, H., Hamdiken, H., Couble, A., Mouniee, D., Boulahrouf, A., et al. (2005). Isolation of actinomycetes producing bioactive substances from water, soil and tree bark samples of the north-east of Algeria. *Journal de Mycologie Médicale*, 15(1), 45-51.
- Kumar, P. S., Ling, C. Y., Zhou, Z. B., Dong, Y. L., Sun, C. L., Song, Y. X., Wong, N. K., & Ju, J. H. (2020). Chemical diversity of metabolites and antibacterial potential of actinomycetes associated with marine invertebrates from intertidal regions of Daya Bay and Nansha Islands. *Microbiology*, 89(4), 483-492.
- Nikolaidis, I., Favini-Stabile, S., & Dessen, A. (2014). Resistance to antibiotics targeted to the bacterial cell wall. *Protein Science*, 23(3), 243-259.
- Pandey, A., Shukla, A. N. U. P. A. M., & Majumdar, S. K. (2005). Utilization of carbon and nitrogen sources by *Streptomyces kanamyceticus* M 27 for the production of an Anti bacterial antibiotic. *African Journal of Biotechnology*, 4(9), 909-910.
- Pandey, B., Ghimire, P., & Agrawal, V. P. (2004). International Conference on the Great Himalaya-yas: Climate, Health, Ecology, Management and Conservation, Kathmandu.
- Pfeifer, Y., Cullik, A., & Witte, W. (2010). Resistance to cephalosporins and carbapenems in Gram-negative bacterial pathogens. *International Journal of Medical Microbiology*, 300(6), 371-379.
- Piddock, L. J. (2012). The crisis of no new antibiotics—What is the way forward? *The Lancet Infectious Diseases*, 12(3), 249-253.
- Quinn, G. A., Banat, A. M., Abdelhameed, A. M., & Banat, I. M. (2020). Streptomyces from traditional medicine: Sources of new innovations in antibiotic discovery. *Journal of Medical Microbiology*, 69(8), 1040-1048.
- Radwan, W. W., AlTuwaijri, D. S., Alwoseamer, A. T., & AlMajed, A. I. (2022). Endodontically treated teeth restoration assessment, decision making and treatment option among dental practitioners in Saudi Arabia. *Annals of Dental Specialty*, 10(1), 118-125.
- Ripa, E. A., Nikkon, K., Zaman, S., & Khondkar, P. (2009). Optimal conditions for antimicrobial metabolites production from a new Streptomyces sp. RUPA-08PR isolated from Bangladeshi soil. *Mycobiology*, 37(3), 211-214.
- Rong, X., Guo, Y., & Huang, Y. (2009). Proposal to reclassify the Streptomyces albidoflavus clade on the basis of multilocus sequence analysis and DNA-DNA hybridization, and taxonomic elucidation of streptomyces griseus subsp. solvifaciens. *Systematic and Applied Microbiology*, 32(5), 314-322.
- Sathi, Z. S., Rahman, M. A. A., & Gafur, M. A. (2001). Identification and in vitro antimicrobial activity of a compound isolated from streptomyces species. *Pakistan Journal of Biological Sciences*, 4, 1523-1525.

- Sebak, M., Saafan, A. E., Abdelghani, S., Bakeer, W., Moawad, A. S., & El-Gendy, A. O. (2021). Isolation and optimized production of putative antimicrobial compounds from Egyptian soil isolate streptomyces sp. MS. 10. *Beni-Suef University Journal of Basic and Applied Sciences*, 10(1), 1-12.
- Shaikh, S., Fatima, J., Shakil, S., Rizvi, S. M. D., & Kamal, M. A. (2015). Antibiotic resistance and extended spectrum beta-lactamases: Types, epidemiology and treatment. *Saudi Journal of Biological Sciences*, 22(1), 90-101.
- Singh, L. S., Baruah, I., & Bora, T. C. (2006). Actinomycetes of Loktak habitat: Isolation and screening for antimicrobial activities. *Biotechnology*, 5(2), 217-221.
- Stanbury, P. F. (2016). *Principles of Fermentation Technology Ed. 3*. Elsevier Science.
- Sujatha, P., Raju, K. B., & Ramana, T. (2005). Studies on a new marine streptomycete BT-408 producing polyketide antibiotic SBR-22 effective against methicillin resistant *Staphylococcus aureus*. *Microbiological Research*, 160(2), 119-126.
- Tiwari, K., & Gupta, R. K. (2012). Rare actinomycetes: A potential storehouse for novel antibiotics. *Critical Reviews in Biotechnology*, 32(2), 108-132.
- Tooke, C. L., Hinchliffe, P., Bragginton, E. C., Colenso, C. K., Hirvonen, V. H., Takebayashi, Y., & Spencer, J. (2019).  $\beta$ -Lactamases and  $\beta$ -Lactamase inhibitors in the 21st century. *Journal of Molecular Biology*, 431(18), 3472-3500. doi:10.1016/j.jmb.2019.04.002
- Vasavada, S. H., Thumar, J. T., & Singh, S. P. (2006). Secretion of a potent antibiotic by salt-tolerant and alkaliphilic actinomycete streptomyces sannanensis strain RJT-1. *Current Science*, 1393-1397.
- Vijayakumar, R., Panneerselvam, K., Muthukumar, C., Thajuddin, N., Panneerselvam, A., & Saravanamuthu, R. (2012). Optimization of antimicrobial production by a marine actinomycete streptomyces afghaniensis VPTS3-1 isolated from palk strait, east coast of India. *Indian Journal of Microbiology*, 52(2), 230-239.
- Yuan, L. J., Zhang, Y. Q., Yu, L. Y., Sun, C. H., Wei, Y. Z., Liu, H. Y., Li, W. J., & Zhang, Y. Q. (2010). Actinopolymorpha cephalotaxi sp. nov., a novel actinomycete isolated from rhizosphere soil of the plant cephalotaxus fortunei. *International Journal of Systematic and Evolutionary Microbiology*, 60(1), 51-54.