

***Nigella Sativa* on the Improvement of Ovarian Dysfunctions in Polycystic Ovary Syndrome (PCOS): A Short Review**

Nurul Hannah Jaafar¹, Sabrina Muhamad Jalani¹, Ruzianisra Mohamed¹, Siti Syairah Mohd Mutalip^{1,2*}

¹Department of Pharmaceutical Life Sciences, Faculty of Pharmacy, Universiti Teknologi MARA, Cawangan Selangor, Kampus Puncak Alam, 42300, Selangor, Malaysia.

²Maternofetal and Embryo Research Group (MatE), Universiti Teknologi MARA (UiTM), 40450 Shah Alam, Selangor, Malaysia.

ABSTRACT

Polycystic ovary syndrome (PCOS) is a multifactorial endocrine disorder that is commonly diagnosed in women of reproductive age. The reported incidents of PCOS are drastically increasing every year worldwide, however, the etiology of PCOS is still not fully comprehended. A traditional herb such as *Nigella sativa* (NS) is being consumed as an alternative to modern medications to manage the signs of PCOS. The effectiveness of NS against PCOS signs and symptoms is being widely published, but studies focusing on its effect on ovarian dysregulations are limited. Hence, this review was written to provide an updated summary of NS and the ovaries of PCOS subjects. This review gathered publications from studies conducted in an in vivo and clinical trial settings and is anticipated to be a reference for future research involving NS and PCOS subjects.

Keywords: Environment, Change, Ecosystem, Destruction, Sustainable development goals (SDGs)

Corresponding author: Siti Syairah Mohd Mutalip

e-mail ✉ syairah@uitm.edu.my

Received: 03 March 2024

Accepted: 18 June 2024

INTRODUCTION

Polycystic ovary syndrome (PCOS)

Polycystic ovary syndrome (PCOS) is one of the most common endocrine disorders in women. It was reported that PCOS had affected around 20% of women globally (Rao *et al.*, 2020). The exact causes of PCOS are still not defined but hyperandrogenism is one of the most reported possible causes of PCOS. Hyperandrogenism results from the imbalance of the hypothalamic-pituitary-ovarian axis (HPO) and oxidative stress (OS) (Ashraf *et al.*, 2019; Sun *et al.*, 2021) which may lead to impaired follicular development (Bertoldo *et al.*, 2019). An imbalance of the HPO axis occurs when there is an increase of gonadotrophin-releasing hormone (GnRH) from the hypothalamus, favoring the production of luteinizing hormone (LH) instead of follicular stimulating hormone (FSH) (Ashraf *et al.*, 2019).

In normal conditions, the development of primordial follicles to primary follicles and subsequently to secondary follicles are gonadotropin-independent while further maturation of preantral follicles to antral follicles is FSH-dependent (Rimon-Dahari *et al.*, 2016; Cox & Takov, 2018). The LH receptors will be formed on the antral follicles as the further development will be LH-dependence. LH will be responsible for stimulating antral follicles into Graafian follicles (Franks & Hardy, 2018). Then, the matured oocyte will be released, and follicles will be

further developed into corpus luteum. In PCOS conditions, the excessive level of androgen will block the development of follicles and a low level of FSH will cause the accumulation of preantral and antral follicles that cannot develop into corpus luteum (**Figure 1**) (John & Michael, 2021). Other than that, many atretic follicles will also be formed due to the excessive amounts of androgen. This is because androgen can go into the layer of granulosa cells of preantral follicles and bind to their cell receptors, resulting in cell death (Zhang *et al.*, 2019; Aburawi *et al.*, 2021).




Figure 1. A normal and polycystic ovary (HealthDirect Australia Limited, 2024)

Nigella sativa – The black seeds

Nigella sativa (NS) is a widely used medicinal plant. It is an annual herbaceous flowering plant (**Figure 2**) from the family of Ranunculaceae and is mostly grown or cultivated in South and Southwest Asia (Ramadan, 2016). Its seeds (**Figure 3**) and oil have been used for centuries to treat various ailments (Tavakkoli *et al.*, 2017), and its efficiency is reported to be due to the different contained constituents. These include terpenes (thymoquinone, dithymoquinone, p-cymene, α -pinene, limonene, carvone, anethol, etc.), steroids (α -hederin), alkaloids (nigellidine, negillidine, nigellidine, nigellidine-N-oxide), flavonoids (quercetin), saponins, fixed oil, coumarins, amino acids, and many others (Rajabian & Hosseinzadeh, 2020; Ahmad *et al.*, 2021). Thymoquinone is the major constituent of the NS oil contributing to its therapeutic potential as an antimicrobial, anti-inflammatory, and antioxidant.



Figure 2. The *Nigella sativa* flowers (Rayhanah Group Sdn Bhd, 2024)



Figure 3. The *Nigella sativa* seeds (Melissa, 2024)

Nigella sativa in PCOS: The reported works

The available reports on the effects of NS on the improvisation of PCOS symptoms are limited. The study reports are mainly on hormonal changes, redox disturbances, histological analysis, and improvement of the menstrual cycle. For instance, a laboratory animal-based study was conducted to observe the protective effect of NS on the number of cystic follicles in letrozole-induced PCOS mice. In this study, both NS seeds powder (10g/kg body weight) and NS oil (4ml/kg body

weight) were used. The results showed that both forms of NS have a similar protective effect which decreased the number of ovarian cystic follicles (Anwar *et al.*, 2016). The same research group also published a study on the ameliorative impact of NS on serum testosterone levels in letrozole-induced PCOS mice, reporting that NS in both liquid and solid states improved serum testosterone levels (Anwar *et al.*, 2021). Other studies on the improvement of NS on body weight (Arivoli, 2019; Nafiu *et al.*, 2019; Ammar & Salem, 2021; Anwar *et al.*, 2021; Naghsh *et al.*, 2023), and the weight, volume, and gross appearance of ovaries in PCOS mice were also reported (Anwar *et al.*, 2021). Supplementation with NS oil to letrozole-induced rats reported a significant increase in the number of rats undergoing regular cycles, the average number of regular cycles, the appearance of corpus luteum, and the reduced number of cystic follicles (Nafiu *et al.*, 2019). This study also reported on the improved levels of circulating gonadotropins, and the increased activities of superoxide dismutase (SOD) and glutathione peroxidase (GPx) (Nafiu *et al.*, 2019). A double-blinded controlled clinical trial conducted on PCOS patients with oligo-amenorrhea, with the intervention of NS oil also indicated that NS has the potential against menstrual irregularities as it shortens the menstrual interval and increases the frequency of the menstrual cycle in PCOS patients (Naeimi *et al.*, 2018; Naeimi *et al.*, 2020). These findings were also supported by a report of a non-randomized clinical trial on 40 PCOS women that showed that a 60-day intake of NS (powder form) resulted in a normal 28-day menstrual cycle in 50% of the women, a normal 3-5 days cycle duration in 85% of the women, and a completely cleared cysts in 15% of the women (Arivoli, 2019). The improving body weight, ovarian morphology, ovulation, and serum LH and FSH following thymoquinone intervention in PCOS-induced Wistar rats were also reported (Javanshir *et al.*, 2018).

The potentials of NS are being continuously reported with the addition of a report on the synergistic effects of NS hydro-alcoholic extract and honey on the hormonal regulations in PCOS-induced rats (Naseran *et al.*, 2020). This laboratory-based study on Wistar rats reported a significant decrease in LH, estrogen, and testosterone; and an increase in FSH level following supplementation with 600mg/kg NS extract and 2400mg/kg honey for 28 days. An increase in the progesterone level also was reported following the intake of 300mg/kg NS extract and 1200mg/kg honey. Similarly, a study by Khani *et al.* (2021) reported that NS hydro-alcoholic extract declined the levels of LH, estrogen, testosterone, and malondialdehyde (MDA) and elevated the levels of progesterone and antioxidant enzymes including SOD, GPx and catalase (CAT) in PCOS-induced rats. NS hydro-alcoholic extract also improved the ovarian histological structure in which the number of Graafian follicles was increased while the number of cystic and atretic follicles decreased (Khani *et al.*, 2021).

Another study (Eini *et al.*, 2020) using NS hydro-alcoholic extract on the quality of oocytes retrieved from PCOS-induced mice during *in vitro* maturation was also reported and indicated that NS extract improves oocyte maturation, oxidative status, and epigenetic modifications. A significant increase was observed in the expression of the *Dnmt1*, *Hdac1*, *Cdk1*, *Mapk*, and *Gpx1* genes with a significant decrease in *Cox2* gene expression of the mature oocytes. These important genes play roles during fertilization and embryonic development

(Eini et al., 2020). Improved ovarian antioxidant activity was also shown through the increase of total antioxidant capacity (TAC) and decline of malondialdehyde (MDA) level following supplementation with NS hydro-alcoholic extract in PCOS-induced rats (Kohzadi et al., 2017).

Combined supplementation of 500mg of metformin and 500mg thymoquinone (NS oil, capsule) showed that NS capsule intake added the improvising effect to metformin as it resumed menstrual regularity, reduced body weight, and regained oxidative balance in the overweight and obese PCOS patients (Jahromi et al., 2023). Another randomized double-blind clinical trial on combined supplementation using NS and black pepper showed an increase in the pregnancy rates, size of dominant follicles, number of follicles, and endometrial thickness compared to the control group which was treated with letrozole and tamoxifen (Jahromi et al., 2023).

The possible effects of Nigella sativa

Various beneficial effects of NS have been suggested as an antimicrobial, cardioprotective, gastroprotective, neuroprotective, anticancer, anti-diabetic, anti-dyslipidemia, anti-obesity, immunomodulatory, anti-inflammatory, nephroprotective, and anti-arthritis activities (Ahmad et al., 2021). NS as an anti-infertility agent has been reported previously, however the particular reports on ovarian function are limited. From the reports discussed above, NS of different forms such as extract, oil, and powder exerted promising effects towards improvisation of the PCOS-induced ovarian dysregulation including folliculogenesis (follicular growth), hormonal levels, structural histology, and oxidative stress. Both experimental studies and clinical trials reported on the enhanced outcomes of the studied parameters, suggesting the promising effects of NS in PCOS patients.

The mechanism of action of NS towards the managing of PCOS-induced reproductive disorders is not fully established, but it was reported to be possibly due to its phytoestrogen effects. Phytoestrogens are derived from plants and possess structures similar to endogenous estradiol (Desmawati & Sulastri, 2019). They can bind to the alpha and beta estrogen receptors (Paterni et al., 2014). Alpha estrogen receptors promote cell proliferation, while beta receptors are responsible for cell apoptosis (Sirotkin & Harrath, 2014). Once the receptor binds to the ligand, it moves from the cytoplasm to the cell nucleus. There, it binds to and influences the area that controls the DNA transcription process or small RNA, which in turn affects the expression of certain genes. Therefore, phytoestrogens have the potential to regulate all processes influenced by estrogen, including the induction of sex hormones (Sirotkin & Harrath, 2014).

Reports on the phytoestrogen effect of NS have been reported earlier in a study conducted to evaluate NS's estrogenic activity using ovariectomized rats (Parhizkar et al., 2011). The study findings showed a significant increase in uterine weight. NS also improved the endometrial histological structure through the increase of luminal and glandular epithelium height, increase of endometrial and myometrial diameters, and increase the number of endometrial glands (Parhizkar et al., 2011). The effect of NS on the uterus was also reported by Sabrina et al. (2023) where co-supplementation of vitamin E and NS resulted in a significantly higher endometrial thickness and the other uterine structures were not disrupted. Other

effects of NS were NS oil reported to help remove uterine content of missed abortion through the reduction of human chorionic gonadotropin (hCG) hormone and facilitation of cervix dilation (Mohammadi et al., 2024). Another review also reported that NS treatment had improved the follicle count, corpus luteum, and level of testosterone and progesterone in infertile women (Darand et al., 2019).

CONCLUSION

The effects of NS supplementation on PCOS-induced ovarian dysfunctions have been continuously reported, despite the molecular mechanism regulating the effects remains unknown. Other than its phytoestrogen effects, the promising effects of NS have been attributed to its potential as an antioxidant, anti-androgenic, anticancer, and anti-inflammatory agent (Ahmad et al., 2013), which is due to the presence of thymoquinone, the primary bioactive compound (Amalia et al., 2022; Alae et al., 2023). The next research focus should be on finding and understanding the mechanism of action of NS, which might require a larger scale of laboratory experiments and clinical trials, as well as determining the optimal dose and duration of intake to achieve effective results.

ACKNOWLEDGMENTS: A special thanks to all staff and members of the Faculty of Pharmacy, UiTM Selangor, Puncak Alam Campus, Malaysia for all the assistance and support provided in completing this study. The authors declare no conflict of interest.

CONFLICT OF INTEREST: None

FINANCIAL SUPPORT: None

ETHICS STATEMENT: None

REFERENCES

- Aburawi, S. M., Treesh, S. A., El Jaafari, H. A., El Ghedamsi, M. T., Nafati, N. A., Benmahmoud, O. A., Almajry, M., & Shebani, N. (2021). Effect of vitamin E on polycystic ovary syndrome induced by dehydroepiandrosterone in female albino mice: Histological study. *Malaysian Journal of Pharmaceutical Sciences*, 19(2), 111-130. doi:10.21315/mjps2021.19.2.8
- Ahmad, A., Husain, A., Mujeeb, M., Khan, S. A., Najmi, A. K., Siddique, N. A., Damanhour, Z. A., & Anwar, F. (2013). A review on therapeutic potential of *Nigella sativa*: A miracle herb. *Asian Pacific Journal of Tropical Biomedicine*, 3(5), 337-352.
- Ahmad, M. F., Ahmad, F. A., Ashraf, S. A., Saad, H. H., Wahab, S., Khan, M. I., Ali, M., Mohan, S., Hakeem, K. R., & Athar, M. T. (2021). An updated knowledge of Black seed (*Nigella sativa* Linn.): Review of phytochemical constituents and pharmacological properties. *Journal of Herbal Medicine*, 25, 100404. doi:10.1016/j.hermed.2020.100404
- Alae, S., Mirani, M., Derakhshan, Z., Koohpeyma, F., & Bakhtari, A. (2023). Thymoquinone improves folliculogenesis, sexual hormones, gene expression of apoptotic markers

- and antioxidant enzymes in polycystic ovary syndrome rat model. *Veterinary Medicine and Science*, 9(1), 290-300. doi:10.1002/vms3.958
- Amalia, A., Hendarto, H., Mustika, A., & Susanti, I. (2022). Effects of nigella sativa on female infertility: A systematic review. In *Proceedings of the 6th International Conference on Medical and Health Informatics* (pp. 234-237). doi:10.1145/3545729.3545776
- Ammar, I. M. M., & Salem, M. A. A. (2021). Amelioration of polycystic ovary syndrome-related disorders by supplementation of thymoquinone and metformin. *Middle East Fertility Society Journal*, 26(1), 1-9. doi:10.1186/s43043-021-00076-1
- Anwar, N. O. R. E. E. N., Nadeem, A. M. E. R., Naheed, K. I. S. H. W. A. R., Jarral, S. A., Abdullah, M. S., Asad, A., & Laique, T. (2021). Impact of Nigella sativa on body weight, volume and weight of ovary in letrozole induced polycystic ovarian syndrome in mice. *Pakistan Journal of Medical & Health Sciences*, 15(9), 2355-2357. doi:10.53350/pjmhs211592355
- Anwar, N., Hamid, S., & Butt, S. A. (2016). Effect of Nigella sativa on several cystic follicles in letrozole-induced polycystic ovaries in mice. *Pakistan Armed Forces Medical Journal*, 66(3), 310-313.
- Anwar, N., Hamid, S., Nadeem, A., Asad, A., Waseem, N., & Mehmood, N. (2021). Effect of nigella sativa on serum testosterone levels in letrozole induced polycystic ovarian syndrome in mice. *Pakistan Armed Forces Medical Journal*, 71(1), 338-342.
- Arivoli, D. (2019). *A Prospective Open Labelled Phase-II Non Randomized Clinical Trial On "Karunjchirakam Chooranam" for Raktha Soorai Vaayu (Poly Cystic Ovarian Syndrome)* (Doctoral dissertation, Government Siddha Medical College, Palayamkottai).
- Ashraf, S., Nabi, M., Rashid, F., & Amin, S. (2019). Hyperandrogenism in polycystic ovarian syndrome and role of CYP gene variants: A review. *Egyptian Journal of Medical Human Genetics*, 20(1), 1-10. doi:10.1186/s43042-019-0031-4
- Bertoldo, M. J., Caldwell, A. S., Riepsamen, A. H., Lin, D., Gonzalez, M. B., Robker, R. L., Ledger, W. L., Gilchrist, R. B., Handelsman, D. J., & Walters, K. A. (2019). A hyperandrogenic environment causes intrinsic defects that are detrimental to follicular dynamics in a PCOS mouse model. *Endocrinology*, 160(3), 699-715.
- Cox, E., & Takov, V. (2018). *Embryology, Ovarian Follicle Development*. StatPearls Publishing LLC, USA.
- Darand, M., Hajizadeh, M., Mirmiran, P., & Mokari-Yamchi, A. (2019). The effect of Nigella sativa on infertility in men and women: A systematic review. *Progress in Nutrition*, 21, 33-41. doi:10.23751/pn.v21i2-S.7088
- Desmawati, D., & Sulastri, D. (2019). Phytoestrogens and their health effect. *Open access Macedonian Journal of Medical Sciences*, 7(3), 495-499. doi:10.3889/oamjms
- Eini, F., Joharchi, K., Kutenaee, M. A., & Mousavi, P. (2020). Improvement in the epigenetic modification and development competence in PCOS mice oocytes by hydro-alcoholic extract of Nigella sativa during in-vitro maturation: An experimental study. *International Journal of Reproductive BioMedicine*, 18(9), 733-746. doi:10.18502/ijrm.v13i9.7668
- Franks, S., & Hardy, K. (2018). Androgen action in the ovary. *Frontiers in Endocrinology*, 9, 452. doi:10.3389/fendo.2018.00452
- HealthDirect Australia Limited. (2024). Polycystic ovarian syndrome (PCOS). Available from: <https://www.healthdirect.gov.au/polycystic-ovarian-syndrome-pcos>.
- Jahromi, R. A., Ansari, A., Zareibabaarabi, Z., Rahmanian, V., & Sharifi, N. (2023). Comparative effect of Nigella sativa+ black pepper and letrozole + tamoxifen on female infertility in women with polycystic ovarian syndrome: A randomized clinical trial. *Journal of Medicinal and Chemical Sciences*, 6(9), 2154-2163. doi:10.26655/JMCHMSCI.2023.9.21
- Javanshir, S. T., Yaghmaei, P., & Hajebrahimi, Z. (2018). Thymoquinone ameliorates some endocrine parameters and histological alteration in a rat model of polycystic ovary syndrome. *International Journal of Reproductive Biomedicine*, 16(4), 275-284. doi:10.29252/ijrm.16.4.275
- John, E. H., & Michael, E. H. (2021). *Female physiology before pregnancy and female hormones*. In: Guyton and Hall Textbook of Medical Physiology. Elsevier, Inc.
- Khani, S., Abdollahi, M., Khalaj, A., Heidari, H., & Zohali, S. (2021). The effect of hydroalcoholic extract of Nigella Sativa seed on dehydroepiandrosterone-induced polycystic ovarian syndrome in rats: An experimental study. *International Journal of Reproductive BioMedicine*, 19(3), 271-282. doi:10.18502/ijrm.v19i3.8575
- Kohzadi, R., Nejati, V., Razi, M., & Najafi, G. (2017). Effects Hydro-alcoholic extract of (Nigella sativa L.) on the level of malondialdehyde (MDA) and total antioxidant capacity (TAC) of the ovary tissue in a rat model of PCOS. *Journal of Animal Environment*, 9(3), 85-92.
- Melissa, P. (2024). Black cumin. Encyclopedia Britannica. Available from: <https://www.britannica.com/plant/black-cumin>
- Mohammadi, B., Nazari Robati, L., Tavakol, Z., & Movahhed, M. (2024). Evaluation of the effect of Nigella sativa oil on the outcome of missed abortion in women: A randomized double-blind clinical trial. *Health Science Reports*, 7(4), e2029. doi:10.1002/hsr.2.2029
- Naeimi, S. A., Hajimehdipoor, H., & Saber, S. (2020). Comparing the effect of Nigella sativa oil soft gel and placebo on oligomenorrhea, amenorrhea and laboratory characteristics in patients with polycystic ovarian syndrome, a randomized clinical trial. *Research Journal of Pharmacognosy*, 7(1), 49-59. doi:10.22127/rjp.2019.206178.1530
- Naeimi, S. A., Tansaz, M., Sohrabvand, F., Hajimehdipoor, H., Nabimeybodi, R., Saber, S., Shakiba, M., & Rohani, M. (2018). Assessing the effect of processed nigella sativa on oligomenorrhea and amenorrhea in patients with polycystic ovarian syndrome: A pilot study. *International Journal of Pharmaceutical Sciences And Research*, 9(11), 4716-4722. doi:10.13040/IJPSR.0975-8232.9
- Nafiu, A. B., Alimi, S., Babalola, A., Ogunlade, A. T., Muhammad, F. D., Abioye, A. R. A. I., Abdulmusawwir, A. O., Oyewole, L. A., Akinola, O., Olajide, O. J., et al. (2019). Anti-androgenic and insulin-sensitizing actions of Nigella sativa oil improve polycystic ovary and associated dyslipidemia

- and redox disturbances. *Journal of Complementary Medicine Research*, 10(4), 186-199. doi:10.5455/jcmr.20190613045154
- Naghsh, N., Moridpour, A. H., Kavyani, Z., Musazadeh, V., Jafarzadeh, J., Safaei, E., Clark, C. C., & Faghfour, A. H. (2023). The effect of *Nigella sativa* (black seed) supplementation on body weight and body composition: A GRADE-assessed systematic review and dose-response meta-analysis of randomized controlled trials. *Journal of Functional Foods*, 105, 105565. doi:10.1016/j.jff.2023.105565
- Naseran, S. N., Mokhtari, M., Abedinzade, M., & Shariati, M. (2020). Evaluation of the effect of *nigella sativa* hydro-alcoholic extract and honey on gonadotropins and sex hormones level in the polycystic ovarian syndrome model of wistar rat. *Scientific Journal of Kurdistan University of Medical Sciences*, 25(1), 117-129. doi:10.52547/sjku.25.1.11
- Parhizkar, S., Latiff, L. A., Rahman, S. A., & Aziz, M. (2011). Evaluation of estrogen-like activity of *Nigella sativa* in ovariectomized rats. *African Journal of Pharmacy and Pharmacology*, 5(8), 1006-1011. doi:10.5897/AJPP10.257
- Paterni, I., Granchi, C., Katzenellenbogen, J. A., & Minutolo, F. (2014). Estrogen receptors alpha (ER α) and beta (ER β): Subtype-selective ligands and clinical potential. *Steroids*, 90, 13-29. doi:10.1016/j.steroids.2014.06.012
- Rajabian, A., & Hosseinzadeh, H. (2020). Dermatological of *Nigella sativa* and its constituent, thymoquinone: A review. In: V.R. Preedy, R.R. Watson. *Nuts and Seeds in Health and Disease Prevention*. Academic Press, Elsevier. doi:10.1016/B978-0-12-818553-7.00024-3
- Ramadan, M. F. (2016). Black cumin (*Nigella sativa*) oils. In *Essential oils in food preservation, flavor and safety* (pp. 269-275). Academic Press.
- Rao, M., Broughton, K. S., & LeMieux, M. J. (2020). Cross-sectional study on the knowledge and prevalence of PCOS at a multiethnic university. *Progress in Preventive Medicine*, e0028. doi:10.1097/pp9.000000000000028
- Rayhanah Group Sdn Bhd. (2024). Kenali Habbatussauda – 3 Fakta Menarik Jintan Hitam. Available from: <https://www.rayhanahmadu.com/blogs/news/kenali-habbatussauda-3-fakta-menarik-jintan-hitam>
- Rimon-Dahari, N., Yerushalmi-Heinemann, L., Alyagor, L., & Dekel, N. (2016). Ovarian folliculogenesis. In: Piprek R. (eds). *Molecular Mechanisms of Cell Differentiation in Gonad Development. Results and Problems in Cell Differentiation*. Springer, Cham. doi:10.1007/978-3-319-31973-5_7
- Sabrina, M. J., Hannah, N. J., Ruzianisra, M., & Syairah, S. M. M. (2023). Co-supplementation of vitamin E & *Nigella sativa* improved the endometrial thickness in mice. *International Journal of Anatomy and Research*, 11(4), 8778-8783. doi:10.16965/ijar.2023.222
- Sirotkin, A. V., & Harrath, A. H. (2014). Phytoestrogens and their effects. *European Journal of Pharmacology*, 741, 230-236. doi:10.1016/j.ejphar.2014.07.057
- Sun, Y., Li, S., Liu, H., Bai, H., Hu, K., Zhang, R., Liu, Q., & Fan, P. (2021). Oxidative stress promotes hyperandrogenism by reducing sex hormone-binding globulin in polycystic ovary syndrome. *Fertility and Sterility*, 116(6), 1641-1650.
- Tavakkoli, A., Mahdian, V., Razavi, B. M., & Hosseinzadeh, H. (2017). Review on clinical trials of black seed (*Nigella sativa*) and its active constituent, thymoquinone. *Journal of Pharmacopuncture*, 20(3), 179-193.
- Zhang, Z., Zhang, Y., Lin, F., Wang, Z., & Zhang, Z. (2019). Endocrine characteristics and regulatory mechanism of follicular development and ovulation failure in the mammalian ovary. In: Wang Z (eds). *Polycystic Ovarian Syndrome*. IntechOpen. doi:10.5772/intechopen.89625