



An Overview on Nasal Polyps' Diagnosis and Management Approach

Ahmed Mahmoud A Alqurashi¹, Saif Abdulaziz H Jawmin¹, Turki Abdulmuin A Althobaiti¹, Mohammad N M F A Aladwani², Abdullah Mohammed E Almuebid³, Joud Fahad Abdullah Alharbi⁴, Mohammed Ali Zarei^{5*}, Lamia Saeed Abdulrahman Al Ghaseb⁵, Waleed Khalid Nawwab⁶, Nedaa Ahmed Zamzamy⁶, Saeed Ali Al Khozaym⁷, Abdulaziz Mohammed Y Hamdi⁸, Muaath Ahmed AlGhamdi⁹

¹Faculty of Medicine, Taif University, Taif, KSA.

²Faculty of Medicine, Jordan University of Science and Technology, Irbid, Jordan.

³Faculty of Medicine, Imam Abdulrahman bin Faisal University, Dammam, KSA.

⁴Faculty of Medicine, Ibn Sina National College of Medicine, Jeddah, KSA.

⁵Faculty of Medicine, King Khalid University, Abha, KSA.

⁶Faculty of Medicine, Umm Al Qura University, Makkah, KSA.

⁷Department of ENT, King Fahad Specialist Hospital, Tabukj, KSA.

⁸Faculty of Medicine, Jazan University, Jazan, KSA.

⁹Department of Family Medicine, Sulaimaniya PHC, Riyadh, KSA.

ABSTRACT

The current management is still controversial and depends upon understanding the complex pathophysiological and etiological factors. There have been several theories that proposed the initial process regarding polyps' formation. Confirming the diagnosis of nasal polyps requires endoscopic investigations or complex radiological investigations. However, currently available medications are mostly based upon clinical experience that describes these pathophysiological and etiological hypotheses. This review provides a wide view on the current evidence-based diagnosis and management abilities in Nasal polyp's conditions, to fully be able to help and aid ENT clinicians to provide good medical compliance throughout the medication process. This review was classified and collected from eligible publicized only English written extracts, articles, and advanced clinical trials. These electronic research engines were included: PubMed and Google Scholar database. Keywords were used together in combination. Treating nasal polyps is directed to relieve symptoms and upgrade therapeutic management to provide the best results of care. The role of endoscopic surgical therapy is highly effective in treating chronic rhinosinusitis if the palliative medical management is found ineffective.

Keywords: Rhinosinusitis, Nasal polyps, Diagnosis, Management, Guidelines

Corresponding author: Mohammed Ali Zarei

e-mail ✉ DR-mozarei@outlook.sa

Received: 19 September 2021

Accepted: 22 December 2021

INTRODUCTION

Nasal polyps present have been a characteristic form of chronic persistent inflammation that is associated with severe and chronic allergic rhinitis hypercreativity reactions, chronic sinusitis, and other chronic inflammatory indeterminate etiologies (Schleimer, 2017; Shamji *et al.*, 2019). Genetical and environmental eosinophilic polyps that are also considered to be responsive and are primarily enhanced sinonasal severity (Stevens *et al.*, 2015; Bochner & Stevens, 2021; Sabbahi, 2022). Confirming the diagnosis of nasal polyps requires endoscopic investigations or complex radiological investigations (Akdis *et al.*, 2015).

The current management is still controversial and depends upon understanding the complex pathophysiological and etiological factors. There have been several theories that proposed the initial process regarding polyps' formation (Cayé-Thomasen *et al.*, 2004; Schleimer, 2017). However, currently available medications are mostly based upon clinical experience that describes these pathophysiological and etiological hypotheses (Ta, 2019; Makhoahle & Gaseitsiwe, 2022). This review provides a wide view on the current evidence-based diagnosis and management abilities in Nasal polyp's conditions, to fully be able to help and aid ENT clinicians to provide good medical compliance throughout the medication process.

MATERIALS AND METHODS

This review was classified and collected from eligible publicized only English written extracts, articles, and advanced clinical

trials. These electronic research engines were included: PubMed and Google Scholar database. The keywords "Nasal" and "Polyps" including words used in MeSH ((('Diagnosis' [Mesh])) , (('Management' [Mesh])) , (('Guidelines' [Mesh])) , (('Treatment' [Mesh])) , (('Biologics' [Mesh])) were used together in integral combinations. This review discussed the first step for diagnosing and treating nasal polyps. All details involving this topic including medical definitions, pathological and physiological classifications, and guidance for treatment were included in this review.

Review

Nasal polyps formation

The different patterns of inflammations in the nasal mucosa have a great impact on the development of nasal polyps in patients with chronic rhinosinusitis issues. Chronic rhinosinusitis is characterized by recurrent nasal congestions, posterior or anterior rhinorrhea, hyposomnia, facial pain for 12

weeks or more (Orlandi et al., 2016; Fokkens et al., 2019; Takabayashi & Schleimer, 2020). Nasal polyps are jelly-cyst-like lesions that originate from the surrounding paranasal sinus cavity and the middle nasal meatus. More intense histological features are found to be edematous pseudo cystic stroma that contains plasma proteins such as albumin and low amounts of collagen production and clear features of fibrosis (Takabayashi et al., 2013; Takabayashi & Schleimer, 2020).

Etiology of nasal polyps

In treating nasal polyps, understanding the causes of these polyps is crucial for achieving effective and appropriate management. Nasal polyps alone are never considered an ultimate diagnosis. The causes of nasal polyps are not fully known or explained and are usually described in endoscopic pathological associations. However, Nasals polyps etiological hypotheses are clarified into several classifications (**Table 1**) (Ta, 2019).

Table 1. Categorical classifications for nasal polyps for explaining nasal polyps' pathological associations: (Ta, 2019)

Anatomical sites of the lesions	Type of polyps	Pathological and physiological etiologies
Localized	Reactive hyperplastic polyps	Inflammatory Neoplastic
Diffused	Host	Anatomical abnormalities Atopic
	Interactive external factors	Staphylococcus superantigen
	Combination of host and external factors interactions	Allergic fungal hypersensitivity rhinosinusitis, eosinophilic fungal rhinosinusitis, and eosinophilic mucin rhinosinusitis.
Systemic	Cystic fibrosis, Churg- Strauss syndrome	

Examination findings and clinical investigations

Since nasal polyps are associated with chronic rhinosinusitis, patients are suspected to present with chronic nasal congestion, decreased or gradual loss of smell, and rhinorrhea (cardinal symptoms of chronic rhinosinusitis). Positive findings are detected using anterior rhinoscopy or other radiological CT

scans [American Academy of Otolaryngology-Head and Neck diagnostic criteria for chronic rhinosinusitis] (**Table 2**) (Rosenfeld et al., 2015). The presence of other comorbid conditions such as asthma and allergic rhinitis may promote the existence of nasal polyps (Feng et al., 2012).

Table 2. Chronic Rhinosinusitis Diagnostic Criteria of the American Academy of Otolaryngology (Rosenfeld et al., 2015).

Symptoms and Signs	Clinical Findings
Nasal Hindrance Nasal effluence Facial discomfort Sleeping disturbances	Physical examination [mucopurulent drainage, polyps, and edema], radiological findings are also preferable.

The investigation for localized nasal polyps must also include microbiological samples, biopsies for further histopathological assessment, particularly for elderly individuals and patients with a high risk of sinonasal malignancies (Feng et al., 2012). Intracranial complications in chronic rhinosinusitis are also suspected and require urgent workup for diagnostic and therapeutic assessment. It is important to refer patients to an otolaryngologist soon after revealing neoplasms, to prevent further complications (Rosenfeld et al., 2015; Sedaghat, 2017).

Treatment and management

The treatment of nasal polyps depends on the different phenotypes and endotypes analyzed to determine the

preferable management [Chronic rhinosinusitis pathophysiology is believed to be associated with T-helper 2 eosinophils, Immunoglobulin E, and high presence of interleukin IL-5, with the dominant presence of cationic eosinophilic protein concentration at the nasal polyp.] (**Table 3**) (Van Zele et al., 2006; Akdis et al., 2015; Ta, 2019). Treating nasal polyps is directed to relieve symptoms and upgrade therapeutic management to provide the best results of care. Since nasal polyps are associated with chronic rhinosinusitis, non-surgical treatment is dedicated to receiving intranasal corticosteroids, performing nasal saline irrigations, receive short- and long-term oral antibiotics to prevent further infections (Rosenfeld et al., 2015; Ajwa et al., 2022).

The main goal of non-surgical treatment is to enhance sinus drainage and improve mucociliary clearance, eliminate local infections and provide a suitable pathway for topical medications. Otolaryngologists should be informed if patients were not responding to medical treatment and other patients

with common comorbidities such as patients who are suffering from asthma or other systemic comorbidities (e.g., Granulomatous diseases, cystic fibrosis, vasculitis, and immunodeficiency) that is involved in the formation of chronic rhinosinusitis (Rosenfeld *et al.*, 2015; Ta, 2019).

Table 3. Phenotypes and endotypes of Nasal polyps in chronic rhinosinusitis: (Ta, 2019)

Phenotypes	Nasal polyps' Mucosal eosinophilia	Nasal polyps and Mucosal neutrophilia	Allergic fungal hypersensitivity rhinosinusitis (AFRS)
Endotypes	Increased eosinophil counts Highly intensive edematous nasal polyps Increased IL-4,13,5 are present Increased macrophages count T-helper 2 (Th2) cells are dominant	High neutrophils count and no eosinophils. Presence of high INF- γ and Negative presence f IL-5 High IL-17 Th1 and Th17 are found dominant	Eosinophil's count is high. the fungal culture is positive Ig-E total is high
Recommended proposed treatment and dosing	Intranasal corticosteroids or oral corticosteroids high doses. leukotriene modifiers Desensitization therapy with Aspirin or salicylate Preferable sinus surgery Biologics: Mepoluziman, Reslizumab, Omalizumab	Intranasal corticosteroid high dose. Sinus surgery Antibiotics; Macrolides	Intranasal or oral corticosteroids Preferable use of antifungals, omalizumab.

The role of endoscopic surgical therapy is highly effective in treating chronic rhinosinusitis if palliative medical management is found ineffective. Endoscopic surgery is done to enhance ventilation and paranasal sinuses drainage, yet enlarge the paranasal pathways for outstanding access for topical medications. However, endoscopic sinuses surgeries are known to improve recurrent symptoms but it doesn't provide a definite cure for the condition, and patients are bound to undergo postoperative medical therapy to maintain recent surgical modifications (Sedaghat, 2017).

CONCLUSION

Nasal polyps present have been a characteristic form of chronic persistent inflammation that is associated with severe and chronic allergic rhinitis hypercreativity reactions, chronic sinusitis, and other chronic inflammatory indeterminate etiologies. Treating nasal polyps is directed to relieve symptoms and upgrade therapeutic management to provide the best results of care. The role of endoscopic surgical therapy is highly effective in treating chronic rhinosinusitis if palliative medical management is found ineffective. The surgical treatment of chronic rhinosinusitis involves endoscopic engagement. However, endoscopic sinuses surgeries are known to improve recurrent symptoms but it doesn't provide a definite cure for the condition, and patients are bound to undergo postoperative medical therapy to maintain recent surgical modifications.

ACKNOWLEDGMENTS: None

CONFLICT OF INTEREST: None

FINANCIAL SUPPORT: None

ETHICS STATEMENT: None

REFERENCES

- Ajwa, N., Faya, I. S. I., Alasbali, S. H. F., Alrayes, A. O. M., Alsaif, A. A., & Aljahdali, S. L. H. (2022). Occurrence of cross-bite among school children in Jeddah City; An observational study. *Annals of Dental Specialty*, 10(1), 59-61.
- Akdis, C. A., Hellings, P. W., & Agache, I. (Eds.). (2015). *Global atlas of allergic rhinitis and chronic rhinosinusitis*. European Academy of Allergic Rhinitis and Chronic Rhinosinusitis.
- Bochner, B. S., & Stevens, W. W. (2021). Biology and function of eosinophils in chronic rhinosinusitis with or without nasal polyps. *Allergy, Asthma & Immunology Research*, 13(1), 8-22.
- Cayé-Thomasen, P., Larsen, K., Tingsgaard, P., & Tos, M. (2004). Basic fibroblast growth factor expression in recurrent versus non-recurrent nasal polyposis. *European Archives of Oto-Rhino-Laryngology and Head & Neck*, 261(6), 321-325.
- Feng, C. H., Miller, M. D., & Simon, R. A. (2012). The united allergic airway: Connections between allergic rhinitis, asthma, and chronic sinusitis. *American Journal of Rhinology & Allergy*, 26(3), 187-190.
- Fokkens, W., Desrosiers, M., Harvey, R., Hopkins, C., Mullol, J., Philpott, C., Alobid, I., Anselmo-Lima, W. T., Bachert, C., Baroody, F., et al. (2019). EPOS2020: Development strategy and goals for the latest European position paper on rhinosinusitis. *Rhinology*, 57(3), 162-168.
- Makhoahle, P., & Gaseitsiwe, T. (2022). Efficacy of disinfectants on common laboratory surface microorganisms at R.S mangaliso hospital, NHLS laboratory, South Africa. *Bulletin of Pioneering Researches of Medical and Clinical Science*, 1(1), 1-12.
- Orlandi, R. R., Kingdom, T. T., Hwang, P. H., Smith, T. L., Alt, J. A., Baroody, F. M., Batra, P. S., Bernal-Sprekelsen, M.,

- Bhattacharyya, N., Chandra, R. K., et al. (2016). International consensus statement on allergy and rhinology: Rhinosinusitis. In *International Forum of Allergy & Rhinology*, 6(S1), S22-S209.
- Rosenfeld, R. M., Piccirillo, J. F., Chandrasekhar, S. S., Brook, I., Ashok Kumar, K., Kramper, M., Orlandi, R. R., Palmer, J. N., Patel, Z. M., Peters, A., et al. (2015). Clinical practice guideline (update): Adult sinusitis. *Otolaryngology-Head and Neck Surgery*, 152(2_suppl), S1-S39.
- Sabbahi, D. A. (2022). Systematic review of different outcomes for dental treatment provided to children under general anesthesia. *Annals of Dental Specialty*, 10(1), 13-33.
- Schleimer, R. P. (2017). Immunopathogenesis of chronic rhinosinusitis and nasal polyposis. *Annual Review of Pathology: Mechanisms of Disease*, 12, 331-357.
- Sedaghat, A. R. (2017). Chronic rhinosinusitis. *American Family Physician*, 96(8), 500-506.
- Shamji, M. H., Thomsen, I., Layhadi, J. A., Kappen, J., Holtappels, G., Sahiner, U., Switzer, A., Durham, S. R., Pabst, O., & Bachert, C. (2019). Broad IgG repertoire in patients with chronic rhinosinusitis with nasal polyps regulates proinflammatory IgE responses. *Journal of Allergy and Clinical Immunology*, 143(6), 2086-2094.
- Stevens, W. W., Ocampo, C. J., Berdnikovs, S., Sakashita, M., Mahdavinia, M., Suh, L., Takabayashi, T., Norton, J. E., Hulse, K. E., Conley, D. B., et al. (2015). Cytokines in chronic rhinosinusitis. Role in eosinophilia and aspirin-exacerbated respiratory disease. *American Journal of Respiratory and Critical Care Medicine*, 192(6), 682-694.
- Ta, N. H. (2019). Will we ever cure nasal polyps? *The Annals of The Royal College of Surgeons of England*, 101(1), 35-39.
- Takabayashi, T., & Schleimer, R. P. (2020). Formation of nasal polyps: The roles of innate type 2 inflammation and deposition of fibrin. *Journal of Allergy and Clinical Immunology*, 145(3), 740-750.
- Takabayashi, T., Kato, A., Peters, A. T., Hulse, K. E., Suh, L. A., Carter, R., Norton, J., Grammer, L. C., Cho, S. H., Tan, B. K., et al. (2013). Excessive fibrin deposition in nasal polyps caused by fibrinolytic impairment through reduction of tissue plasminogen activator expression. *American Journal of Respiratory and Critical Care Medicine*, 187(1), 49-57.
- Van Zele, T., Claeys, S., Gevaert, P., Van Maele, G. E. O. R. G. E. S., Holtappels, G., Van Cauwenberge, P., & Bachert, C. (2006). Differentiation of chronic sinus diseases by measurement of inflammatory mediators. *Allergy*, 61(11), 1280-1289.