



## Breeding ecology of the common coot (*Fulica atra*) at El-Golea Lake (Algerian Sahara)

Guergueb El-Yamine<sup>1-3</sup>, Nouidjem Yassine<sup>2-3\*</sup>, Bounab Choayb<sup>1-3</sup>,  
Bensaci Ettayib<sup>2-3</sup>, Haddad Soumia<sup>1-3</sup>, Houhamdi Moussa<sup>3</sup>

<sup>1</sup> Department of Biology, University of Ghardaia, Algeria

<sup>2</sup> Department of Natural and Life Sciences, University of M'Sila, Algeria

<sup>3</sup> Biology, Water and Environment Laboratory (LBEE), University 8 Mai 1945 Guelma, BP 401, Guelma, 24000, Algeria

### ABSTRACT

The breeding ecology of common coot *Fulica atra* was investigated at El-Golea Lake, in the northern part of Algerian Sahara, where the breeding season was comparatively short, during three months. The distinct habitats within the lake used by common coots for nesting are associated with *Typha angustifolia* (96%) and *Scirpus lacustris* (04%), the vegetation covers the density that varies between 30% to 80%. The mean clutch size was  $5.6 \pm 3.2$  eggs ( $N = 45$  clutches) and the laying occurred mainly in April, but continued until the end of May. The breeding success rate was (75.55%) with the predation rate of (72.72%), and clutch desertion of (27.27%). The effects of disturbance on the breeding success of this water bird and the necessity to protect this species and also this wetland were discussed in the present study.

**Keywords:** Breeding success, *Fulica atra*, El-Golea Lake, Sahara

**Corresponding author:** Yassine Nouidjem

**Received:** 25 September 2017

**Accepted:** 08 January 2018

Haouam et al., 2006; Zitouni et al., 2013), the Hauts Plateaux (Samraoui & Samraoui, 2007; Baaziz & Samraoui, 2008), and Algiers region (Metna, 2014 et al.; Metna et al., 2015). The present study focused on the breeding ecology of this species in the northern Algerian Sahara at El-Golea Lake.

### INTRODUCTION

The common coot *Fulica atra* is one of the most abundant water birds in Europe (Jortay, 2002), in North Africa (Isenman & Moali, 2000), and in Algeria (Baaziz & Samraoui, 2008; Metna et al., 2013) which range includes all of Northern Algeria including the Hauts Plateaux (Ledant et al., 1981; Isenmann & Moali, 2000; Metna et al., 2013; Guergueb et al., 2014) that presents nesting areas in southern Algeria at some oasis (Etchecopar & Hue, 1964).

This water bird is the common species of the Algerian wetlands (Rizi et al., 1999). It has a Breeding sedentary status in all the Mediterranean basin (Allouche & Tamisier, 1984; Samraoui & Samraoui, 2007). In Algeria, the reproductive biology of the common coot *Fulica atra* was studied in some areas of the country, particularly in Numidia region (Rizi et al., 1999;

### MATERIAL AND METHODS

#### Study area

The lake El-Golea (30° 31 'N, 02° 56.'E), a salt marsh has a surface of 270 km<sup>2</sup> for a maximum depth of 02.5 m located in the northern Sahara (D.G.F, 2005). This region has an arid Mediterranean climate, characterized by hot and dry summer, the average annual rainfall is 20 mm. Between the Grand Erg Oriental in the East and Grand Erg Occidental in the west, this lake composed of two interconnected water bodies, an upper basin which is a permanent brackish water lake, very rich in biodiversity, fed mainly by wastewater from the city of Menea and Hassi El-Gara, and a lower basin very broad, composed of saline deposits (Khalifa et al., 2009). (Fig 01).




Figure 01. Map of the study area.

**Data collection and statistical analysis**

This work consists of studying and characterizing reproduction of the common coot *Fulica atra*. It has been considered between January 2014 to June 2014 at El-Golea Lake. For this, a nest monitoring was carried out during the entire breeding season twice a week. The date of nest establishment, the start of egg laying, egg size, egg biometrics (large and small diameter) and nest characteristics (Composition, internal and external diameter, depth of nests and their height relative to water) were the parameters which were taken into consideration. Nests and eggs have been marked to better track lying from the start, and the measurements were made with an electronic caliper at 1 / 10th of a mm. The vegetation was grouped into four classes: *Scirpus lacustris*, mixed *S. lacustris*, mixed *Typha angustifolia* and other vegetation.

**RESULTS AND DISCUSSION**

**Breeding habitats' characteristics**

During the research, 45 nests were identified in the upper basin of the body of water during the whole study period, which spanned a breeding season from February to June in 2015 (03 nests during the month of February, 08 nests during the month of March, 21 nests in the month of April, 11 nests in the month of May, and 02 nests in the month of June).

Most of the nests were associated with *Typha angustifolia* (96%), and *Scirpus lacustris* (04%), with a vegetation cover density that varies between 30% to 80%, and mean water

depth of  $68,8 \pm 17,55$  (Table 01). These nests have an average outside diameter of  $30,15 \pm 2,98$  cm with a maximum of 35 cm, an average internal diameter of  $17,23 \pm 1,83$  cm with a maximum of 20 cm, and an average depth of  $3,81 \pm 1,65$  cm. The nests were positioned at an average height of  $13,38 \pm 1,92$  cm in relation to the water. The breeding strategy of this water bird is related to the density of the vegetation strata (*Phragmites australis*) (Huhta *et al*, 1998).





Figure 02. Percentage of vegetation use by nesting coots at El-Golea Lake

**Table 01.** Nest characteristics and egg measurements of common coot at El-Golea Lake

	Mean ± SEM	Max.	Min.	N
<b>Nest characteristics</b>				
Distance to shore (m)	48,82±36,98	100	4	45
External diameter (cm)	30,15±2,98	35	20	45
Internal diameter (cm)	17,29±1,83	22	14	45
Nest depth (cm)	3,91±1,65	7	2	45
Nest height (cm)	13,38±1,92	17	10	45
Water depth (cm)	68,8±17,55	99	40	45
Vegetation height (cm)	118,66±39,11	190	25	45
Vegetation density (%)	48,89±15,88	80	20	45
<b>Egg measurements</b>				
Egg length (mm)	54,13±1,67	58,36	49,52	257
Egg breadth (mm)	36,71±0,84	38,81	34,37	257

**Clutch size**

During the study, it was observed that female common coots lay between 1 and 11 eggs per clutch with an average of 5.6 ± 3.20 eggs per nest. 24.44% of broods (11 broods) with 6 eggs, 28.89% of broods (13 broods) with 7 eggs, 15.55% of broods (7 broods) with 5 eggs, 8.89% broods (4 broods) with 4 eggs, and 6.67% broods (3 broods) with 3 eggs, were found. Only 4.44% broods (2 broods) with 1 and 8 eggs, and 2.22% broods (1 brood) with 2, 9 and 11 eggs, were found.




**Figure 03.** Frequency distribution of clutch size of coots.

**Table 02.** Egg measurements data in different studies of the breeding ecology of common coots in Algeria

	Clutch size Average (Min - Max)	Dates (Start - Finish)
Tonga Lake in 1997 by (Rizi et al., 1999)	4,15 (1 - 8)	/
Timerganine Lake in 2005 by (Samraoui & Samraoui, 2007)	7,2 (1 - 16)	(18 March - 02 July)
Réghaia Lake in 2010 by (Metna et al., 2013)	4,8 (1 - 10)	(10 April - 10 May)
This study	5,6 (1 - 11)	(20 February - 12 June)

**Egg-Laying period and laying size**

Egg-laying by common coots started on February 20th and ended on June 12th, the mean egg-laying date was on April 17th. The laying size was between 1 and 11 eggs with an average of 5.6 ± 3.2 eggs. The results of this study are more important than those found in the coastal areas (Tonga Lake by Rizi et al., 1999 and Lake Reghaia by Metna et al., (2013), but are inferior to those cited by Samraoui and Samraoui (2007) in the Timerganine site, that may be due to the phenomenon of parasitism or the age of females (Table 02) according to Samraoui and Samraoui (2007). This phenomenon is well known in the region of Numidia.



**Figure 04.** Distribution of laying period of coots at Lake El-Golea

**Egg measurement**

The 45 surveyed nests contained a total of 257 eggs. Figure 04 summarizes all the information on the characteristics of the eggs including the dimensions, the volumes and their shape indices. The large diameter of the 257 eggs measured was on an average of 54.14 ± 1.67mm with the maximum of 58.36mm, and the minimum of 49.52mm; the small diameter of the eggs was between 36.71 ± 0.84mm with the maximum of 38.81mm, and the minimum of 34.37mm. The lowest form factor of the 257 eggs studied was 17.87mm, and the strongest was 22.19mm, which gave us an average of 19.87 ± 0.84. The calculated volumes of the eggs were on an average of 34.75 ± 2.10 mm<sup>3</sup>.

The large diameter of the eggs measured in El-Golea Lake varied from 49.52 to 58.36 mm, giving us an average of 54.13 ± 1.67 mm. The small diameter varied from 34.37 to 38.81 mm with an average of 36.71 ± 0.84 mm. These results were similar to those found by Samraoui and Samraoui (2007) in the Timerganine wetland, and those found by Metna et al. (2013) in the Reghaia wetland and by Rizi et al. (1999) in the Tonga Wetland (Table 03).

The results were also similar to those found by several authors in Europe, which meant that this study site offers all the necessary conditions, especially the food necessary for the good development of the eggs of coots.

**Table 03.** Nests' size data in different studies of the breeding ecology of common coots in Algeria

	Small diameter (mm) Moy ± Standard deviation (Min - Max)	Grand diamètre (mm) Moy ± Standard deviation (Min - Max)
Tonga Lake in 1997 by (Rizi et al., 1999)	35 ± 0,7 (33,2 - 36,9)	52,4 ± 1,9 (48,4 - 58,2)
Timerganine Lake in 2005 by (Samraoui &	36,4 ± 1 (33 - 40,4)	53,1 ± 0,1 (42,5 - 59,8)

Samraoui, 2007)		
Réghaia Lake in 2010 by (Metna et al., 2013)	37,3 ± 1,3 (31,8 - 40)	53,5 ± 2,3 (44,6 - 59,5)
This study	36,8 ± 0,9 (34,1 - 38,8)	54,2 ± 1,7 (49,5 - 58,4)

#### DISTURBANCE

This study highlighted the strong anthropogenic pressure on coots or other species at El-Golea Lake. Human disturbance was recognized as a threatening process that can affect habitat use or lower the breeding success of water birds (Flemming et al. 1988; Anderson and Keith 1980). The disturbance had an impact during those periods of the annual cycle when the species was most vulnerable to high energy demands like incubation.

#### CONCLUSION

This study presented new data on the reproductive ecology of the common coot *Fulica atra* at El-Golea Lake (Algerian Sahara); it is the first study in this region which indicated that the species is a regular breeder in the area, and also confirmed the reproduction of this species in wetlands of Algerian Sahara and their important role in its generation. The breeding strategy of this water bird is related to the density of the vegetation strata (*Phragmites australis*). Nouri et al (2013) suggested that the coots evaluate the level of the nest depredation before start to breed. In addition, the good nesting adaptive strategy of this common bird is influenced by the anthropic disturbance.

The cycle of reproduction of the common coot *Fulica atra* is faced by many troubles that can affect the breeding success (the habitat destruction and the excessive disturbance). This result can give us an image on the stability of the reproduction of this species despite the severe conditions. For this reason, some actions which are necessary to protect this species and also these wetlands should be carried out.

#### REFERENCES

1. ALLOUCHE, L. et TAMISIER, A. (1984). - Feeding convergence of Gadwall, Coot and the other herbivorous waterfowl species wintering in the Camargue. *Wildfowl*, 35, 135-142.
2. Anderson DW, Keith JO (1980) The human influence on seabird nesting success: conservation implications. *Biological Conservation* 18: 65-80.
3. Baaziz N, Samraoui B (2008) Status and diurnal behaviour of wintering common coot *Fulica atra* L. in the Hauts Plateaux, Northeast Algeria. *Eur. J. Sci. Res.*, 23 (3), 495-512.
4. D.G.F (2005) Ramsar wetlands fact sheet «Sebkhet El Melah» (General Directorate of Forests - Wilaya of Ghardaia) », 13 p.
5. Etchecopar RD, Hüe F (1964) *The Birds of North Africa*. Edition Boubée & Co, Paris. 606 p.
6. Flemming SP, Chiasson RD, Smith PC, Austin-Smith PJ, Bancroft RP (1988) Piping Plover status in Nova Scotia related to its reproductive and behavioural responses to human disturbance. *Journal of Field Ornithology* 59: 321-330.
7. Guergueb EY, Bensaci E, Nouidjem Y, Zoubiri A, Kerfouf A, Houhamdi M (2014) Overview of the waterbird diversity of Chott El-Hodna (Algeria). *Bull. Soc. zool. Fr.*, 2014, 139 (1-4): 233-244.
8. Haouam L, Samraoui F, Samraoui, B (2006) Influence trophiques resources on the reproduction of the common coot *Fulica atra* in the wetlands

- Numidie eco-complex (Algeria). *Bull. INSTM, Salambo*, p. 17-20.
9. Huhta E, Jukka J, Pekka R (1998) Distribution and reproductive success of the Pied Flycatcher *Ficedula hypoleuca* in relation to forest patch size and vegetation characteristics; the effect of scale." *Ibis* 140.2: 214-222.
  10. Isenmann P, Moali A (2000) *Birds of Algeria / Birds of Algeria*. Paris: Ornithological Studies Society of France.
  11. Jortay A (2002) Nesting and wintering the common coot (*Fulica atra*) in the settling ponds of Hesbaye liégeoise, *Aves*, 39/2.
  12. Khelifa R, Youcefi A, Bouzid A, Bouchecker A, Boukhssaim M, Samraoui B (2009) A new breeding site for the Greater Flamingo *Phoenicopterus roseus* in Algeria, *Flamingo* 17.
  13. Ledant JP, Jacob JP, Jacob P, Malher F, Ochando B, Roche J (1981). Update of the Algerian avifauna. *The Gerfaut* 71; 295 - 398.
  14. Metna F (2014) *Ethology and nesting biology of Ferruginous Duck Aythya nyroca (Guldenstadt, 1770) and pochard Aythya ferina (Linnaeus, 1758) in the nature reserve of Réghaia Lake.*, Phd thesis, University of Tizi Ouzou (Algeria), 140p.
  15. Metna F, Lardjane-Hamiti A, Boukhemza-Zemmouri N, Boukhemza M, Merabet S, Abba R (2015) Diet of the Coot *Fulica atra* (Aves, Rallidae) in the nature reserve of Lake Réghaia (Algiers, Algeria), *Zoology and Ecology*, DOI: 10.1080/21658005.2014.994363.
  16. Metna F, Lardjane-Hamiti A, Merabet S, Sayoud MS, Boukhemza-Zemmouri N, Boukhemza M (2013) Monthly changes in the workforce, characteristics of the nests and eggs of the common coot (Aves, Rallidae) in the nature reserve of Lake Réghaia (Algeria), *Bull. Soc. zool. Fr.*, 2013, 138(1-4): 93-101.
  17. Nouri N, Guercueb E, Merabet O, Bouslama Z, Houhamdi M (2013) Ecological Characteristics of breeding common coot *Fulica atra* (Rallidae) in the Central Highlands of Constantine (East of Algeria), *European journal of Scientific research*, 2013, 99 (3): 362-373.
  18. Rizi H, Benyacoub S, Chabi Y, Ban'bura J (1999) Nesting and reproductive characteristics of coots *Fulica atra* breeding on two lakes in Algeria, *Ardeola* 46: 179-186.
  19. Samraoui F, Samraoui B (2007) The reproductive ecology of the Coot *Fulica atra* L. in the Hauts Plateaux, northeast Algeria. *Waterbirds*, 30, 133-139.
  20. Zitouni A, Rouibi AH, Bara M, Tahar A, Houhamdi M (2013) The common coot *Fulica Atra* in the Northeast of Algeria (National park of El Kala): study of the breeding biology, *Scholars Research Library*, 4 (10):92-95.