



## Larvicidal Efficacy of Mentha Pulegium Essential Oil Against Culex Pipiens L. And Aedes Caspius P. Larvae

Radja Guenez<sup>1,2</sup>, Fouzia Tine-Djebbar<sup>1,2\*</sup>, Samir Tine<sup>1,2</sup>, Noureddine Soltani<sup>2</sup>

<sup>1</sup>Department of Nature and Life Sciences, Larbi Tebessi University, Tebessa, Algeria.

<sup>2</sup>Laboratory of Applied Animal Biology, University of Badji Mokhtar, Annaba, Algeria

### ABSTRACT

Mosquitoes are generally controlled by conventional insecticides which poses strong secondary effects on the environment. In this context, the present study examines the chemical composition of the essential oil of *Mentha pulegium* (Lamiaceae) and its Larvicidal activity against the two mosquito species *Culex pipiens* and *Aedes caspius*.

*Culex pipiens* and *Aedes caspius* larvae were collected in the year 2015 from untreated areas located at Tébéssa (Northeast Algeria). Aerial parts of *Mentha pulegium* were harvested during April 2015 from the Tebessa area. After the dryness, the plant material was subjected to hydrodistillation using a Clevenger-type apparatus. Chemical composition of the essential oil was determined using a gas chromatography-mass spectrometry analysis. Toxicity test was made on the fourth-instar larvae according to the World Health Organization recommendations. Gas chromatography-mass spectrometry analysis led to the identification of 14 components. Pulegone (72.50%), Eucalyptol (10.44%) and P-menthone 2-ethyl-5-propyle (5.46%), are the major constituents. Biological test revealed that the *M. pulegium* essential oil exhibits a larvicidal activity and *A. caspius* is more sensitive than *C. pipiens* larvae.

*M. pulegium* essential oils, constitute an alternative to the conventional insecticides for controlling mosquitoes.

**Keywords:** Mosquitoes, *Mentha pulegium*, CPG/SM, Chemical composition, Toxicity.

**Corresponding author:** Fouzia Tine-Djebbar

**e-mail** ✉ [fouzia.djebbar@yahoo.fr](mailto:fouzia.djebbar@yahoo.fr)

**Received:** 12 September 2017

**Accepted:** 18 December 2017

### 1. INTRODUCTION

Mosquitoes can transmit serious human diseases such as malaria, dengue, filariasis, and yellow fever, which affect more than 700 million people annually throughout the world (Schaffner et al. 2001; Ghosh et al. 2012). To prevent the proliferation of this arthropod and to improve the environmental quality and public health, more attention has been focused on botanicals, which are effective, eco-friendly, biodegradable and inexpensive, and one of the possible alternatives to synthetic insecticides (Choochate et al. 2005). Previous studies have revealed larvicidal, pupicidal and repellent activities of plant extracts against mosquitos (Shaalani et al. 2005; Pavela et al. 2014; Rocha et al. 2015; El Akhal et al. 2015; El Akhal et al. 2016). More than 2000 plant species have been known to produce secondary metabolites involved in pest control programs (Ghosh et al. 2012). The members of the plant families, Myrtaceae, Asteraceae, Lamiaceae, Miliaceae, Rutaceae, Cupressaceae, Zingiberaceae and Apiaceae, have been examined for anti-insect activities (Onyambu et al. 2015). Essential oils provide a rich source of biologically active Monoterpenes and are well documented for the bioactivities against the insect pests (Govindarajan, 2010). Some of the essential oils with a promising mosquito control potential are plants from genus *Tagetes* spp (Dharmagadda et al. 2005), *Ocimum* spp (Bhatnagar et al. 1993), *Cymbopogon* spp (Ansari and Razdan, 1995), *Mentha* spp (Ansari et al. 2000), *Thymus vulgaris* (Bouguerra et al. 2017; Dahchar et al. 2016). *Mentha* is

an important genus of Lamiaceae, which recognized for their vital oils, medicinal uses and antimicrobial activity of different species (Skaltsa et al. 2003). It consists of about 25–30 species and most of them are grown in the temperate regions of Australia, South Africa and the Eurasia region. *Mentha pulegium* L. is a native of temperate Asia, Africa, and Europe (G.R.I.N, 2010). The previous studies have reported that the various *Mentha* spp. plant extracts exhibited the Larvicidal activity against several mosquito species such as *C. pipiens*, *C. quinquefasciatus*, *A. aegypti*, *Anopheles stephensi* and *An. tessellatus* (Amer and Mehlhorn 2006; Koliopoulos et al. 2010; Samarasekera et al. 2008; Sukumar et al. 1991; Traboulsi et al. 2002; Ambindei et al. 2017). *C. pipiens* and *A. caspius* are the most interesting mosquito species in Algeria, particularly in the Tebessa area (Tine-Djebbar et al. 2016). This present study aims 1/ to determine the chemical composition of the essential oils of cultivated mint species largely used in Algeria. *M. pulegium* and 2/ to test their insecticidal activities against *C. pipiens* and *A. caspius*.

### 2. MATERIALS AND METHODS

#### 2.1. Mosquito rearing

*Culex pipiens* Linnaeus, 1758 and *Aedes caspius* Pallas, 1771 larvae were collected in the year 2015 from untreated areas located at Tebessa (Northeast Algeria). The larvae specimens were morphologically identified according to the identification keys of Brunhes et al. (2000) and Himmi et al. (1995). The pyrex storage jars (80 by 100mm) containing 150 ml of tap water were maintained at a 25 °C temperature and a photoperiod of 14:10 (L:D). The larvae were daily fed with fresh food consisting of a mixture of Biscuit Petit Regal-dried yeast (75:25 by weight) and water was replaced every four days.

## 2.2. Plant Material

The Aerial parts of *Mentha pulegium* Linnaeus, 1753 were harvested during April 2015 from the Tebessa area (Northeast Algeria). Then, the plant parts were washed with tap water, to eliminate soil and other surface contaminants. After the dryness, at laboratory temperature and obscurity, the plant material was cut into small pieces.

## 2.3. Extraction of the essential oils

100g of the air-dried aerial parts of *Mentha pulegium* were subjected to hydrodistillation for 3 hours with 500ml distilled water using a Clevenger-type apparatus. The oil obtained was collected and dried over anhydrous sodium sulfate and stored in a screw capped glass vials in a refrigerator at 4 °C prior to the analysis. The yield based on the dried weight of the samples was calculated (Bouguerra et al. 2017).

## 2.4. Gas chromatography-mass spectrometry

The essential oil of *M. pulegium* was subjected to a gas chromatography-mass spectrometry (GC-MS) analysis using the Perkin Elmer chromatography Autosystem GC. The column HP-5 (length: 30m with an internal diameter of 0.25mm and a film thickness of 0.25 $\mu$ m) was used with helium as a carrier gas. The GC oven temperature was kept at 60 °C for 1 minute and programmed to 200 °C for 13 minutes. The injector's temperature was set at 250 °C. The sample was dissolved in acetone. A volume of 1 $\mu$ l was injected for the GC-MS analysis. The essential oil was analyzed in the Laboratory of Pharmacognosy (Department of Pharmacy, Faculty of Medicine, Batna).

## 2.5. Toxicity Bioassay

Bioassays were conducted as previously described (Boudjelida et al. 2005). The essential oil of *M. pulegium* was added to the treatment beakers at different final concentrations. The newly ecdysed fourth-instar larvae of *C. pipiens* and *A. caspius* (< 8 hours old) were exposed to the different concentrations (50, 100, 250, 500, and 1000 ppm) for 24hours in accordance with the World Health Organization (WHO) criteria (Anonym 1983). The positive controls were exposed to ethanol (solvent) while the negative controls were exposed to water only. Mortality was registered after 24hours of treatment. The test was carried out with 4 replicates containing 25 larvae each per concentration. The mortality percentages obtained were corrected (Abbott 1925) and the toxicity data subjected to the probity analysis (Finney 1971). The Lethal concentrations (LC<sub>50</sub> and LC<sub>90</sub>) and 95% confidence limits (95% CL) were estimated, and the slope of the concentration-mortality lines was calculated (Swaroop et al. 1966).

## 2.6. Statistical Analysis

The number of individuals tested in each series is given with the results. The data are presented as the mean  $\pm$  standard deviation (SD). The significance of the different series was tested using the Student's t test. All the statistical analyses were performed using the MINITAB Software (Version 17, PA, State College, USA) and  $p \leq 0.05$  was considered to be a statistically significant difference.

## 3. RESULTS AND DISCUSSION

### 3.1. Extraction yield

From the results of steam distillation, we found that the yield of extraction of the essential oil of *M. pulegium* was  $0.87 \pm 0.055$  % of the dry matter of the aerial part of the plant. This yield is expressed in milliliters for 100g of dry matter. The obtained yield in our experiments is lower than the EOs extracted from the same species collected in Lavras, Brazil (2.54%) (Silva et al.

2015), in M'irt (Morocco) (5.2%) (Amalich et al. 2016) and in Reguiba, El-Oued (South-East Algeria) (2.34%) (Ouakouak et al. 2015). Compared to the yield of pennyroyal harvested in Moroccan North (1.9%) (Cherrat et al. 2014) and the one obtained by Boughdad et al. (2011), the yield obtained in this study is also higher. It is relatively comparable to other plants industrially exploited as a source of essential oils like *Artemisia herba alba* (0.59%) and *Artemisia absinthium* (0.57%) (Derwich et al. 2009), *Thymus vulgaris* (1%) (Imelouane et al. 2009), *Mentha pulegium* and *M. spicata* (1% and 0.87%, respectively) (Boukhebt et al. 2011). In contrast, Benayad (2008) and Mahboubi and Haghi (2008) reported in the same plant a yield of 2.33% and 0.27 % respectively. However, the EO yield obtained from 7 species of the genus *Mentha* ranged from 0.69% to 2.33% (Pavela et al. 2014) and the total yield for *L. stoechas* is around 2.17% (Ben Slimane et al. 2015). Zantar et al. (2015) show a greatest yield for *M. pulegium* (3.5%) and *T. vulgaris* (3.6%). The high yield of EOs obtained during the full-flowering period, is probably due to its ecological role in attracting pollinators and in being an antifungal defense mechanism (Verma et al. 2010). These percentages were similar to those reported by Hmiri et al. (2011). In addition, the difference in the yield of the *Mentha* essential oils with respect to the geographical regions was reported (Abdullah, 2009).

### 3.2. Chemical composition of essential oil

The percentages and the retention times of the identified compounds of EOs of *M. pulegium* are listed in Table 1 and Figure 1. Fourteen compounds, representing 99.29% of the total essential oil, were identified. The major component was Pulegone (72.50%), other components present in the appreciable contents were: Eucalyptol (10.44%), Menthone (5.46%), Pentanone (3.78%), L-  $\beta$  Pinene (1.90%), Piperitenone (1.51%) and Isopulegone (1.12%).

The literature data suggest that *M. pulegium* is a chemical Polymorph species in both the qualitative and/or quantitative composition (Kokkini et al. 2004). Compared to the published data, the chemical profile obtained presented differences, but also some similarities. 38 components were identified in the essential oils of *M. pulegium* collected from Tunisia (Hafedh et al. 2009) and Iran (Sardashti and Adhami 2013) and 16 from Morocco (Benayad 2008). It has been found that the *M. pulegium* L. oils from Bulgaria contains major components, Pulegone (42.9–45.4%) (Stoyanova et al. 2005); from Uruguay; pulegone (73.4%), isomenthone (12.9%) (Lorenzo et al. 2002); from Egypt; Pulegone (43.5%), Piperitone (12.2%) (El-Ghorab, 2006); from Tunisia, Pulegone (41.8%), Isomenthone (11.3%) (Mkaddem et al. 2007). Boukhebt et al. (2011) studies showed three chemotypes of *M. pulegium* with the following major oil components: Pulegone, Piperitenone, and Isomenthone.

The chemical composition of the *Mentha pulegium* L. oil has been subjected to numbers of studies, which have shown a difference in its constituents depending on the region of cultivation and there have been some variations in the constituents from the different countries (Boukhebt et al. 2011). Studies reported by Mahboubi and Haghi (2008) show clearly that the *M. pulegium* essential oil contains the following compounds: Piperitenone 33.0%,  $\alpha$ -terpineol 4.7%, Pulegone 2.3% from which Piperitone was the major constituent (38.00 %). In contrast, Pulegone (88 %) was the main component in the same plant analysed by Ouraini et al. (2005). In addition, the results from Vian et al. (2008) present that Pulegone (83.70 %) was the main component of the *M. pulegium* essential oil, whereas, Hilan et al. (2007) found that menthol is the main constituent (50%). The essential oils of *M. spicata* and *M. pulegium* mainly consisted of oxygenated monoterpenes and

carvone, menthol and menthone were the principal constituent (Boukhebtli et al. 2011).

Several reports revealed that the essential oil contents depend not only on the temperature and relative humidity but also on the duration of sunshine, air movement, and rainfall (Kastner 1969; Vernet et al. 1977). In the leaf essential oil *Mentha spicata*, 57 compounds and carvone (59.40%) is the major component, with other components which was present in appreciable contents such as: limonene (6.12%), 1,8-cineol, germacrene-D (4.66%),  $\beta$ -caryophyllene (2.969%),  $\beta$ -bourbonene (2.796%),  $\alpha$ -terpineol (1.986%) and terpinene-4-ol (1.120%) (Boukhebtli et al. 2011). In other studies (Soković et al. 2009), carvone (49.5%) is also the major constituent, followed by menthone (21.9%) and limonene (5.8%) of the extractable essential oils from this plant species.

The GC/MS analysis of the *Lavandula stoechas* essential oil led to the identification of 20 components. Camphor (36.14%), 1,8-Cineole (25.16%), camphene (11.44%) and fenchone (9.08%), were the major constituents of which (El Ouali Lalami et al. 2016) 29 components representing 96.43% of the essential oil extracted from *Mentha spicata* cultivated at Ghardaia were identified with cis-carvone oxide, 1,8-cineole, cis-dihydrocarvone, and limonene (5.80%) as the main components (Laggoun 2016). Various factors as the environmental conditions, the extraction technique, the drying, the period and gathering sites, the plant age (Aberchane et al. 2001; Bourkhiss et al. 2009; Okoh et al. 2007; Bourkhiss et al. 2011), the concentration of the extract and the concentration of its active components, can influence the performance and the chemical composition of the extract.

### 3.3. Larvicidal Activity

In the present study, the dose-response relationship was determined for the *M. pulegium* essential oil applied to a newly ecdysed fourth instar larvae of *C. pipiens* and *A. caspius* (Table 2). The mortality was scored at 24 hours after treatment. With probit,  $LC_{50}$  was calculated as 38.75 and 28.16ppm, and  $LC_{90}$  was 85.91 and 53.75ppm for *C. pipiens* and *A. caspius*, respectively (Table 3). *A. caspius* is the most sensitive species. After the treatment, the intoxicated larvae showed a change in their behaviour by sinking to the bottom of the jar and remain there motionless until they died. Our results indicate that the *M. pulegium* essential oils and their active components could be developed as control agents against the mosquito larvae.

The efficacy of the botanical extracts against the population of mosquitoes can vary depending on the plant species, age and parts of the plant, the physicochemical characteristics and chemical composition of EO, the environmental conditions, the used extraction technique, the drying process, the period and growing environment and the cultivation practices (Sukumar et al. 1991; El Ouali Lalami et al. 2016). The insecticidal properties of EO are very well documented (Traboulsi et al. 2002; Koul et al. 2008; Pavela, 2008a; Pavela, 2008b; Urzua et al. 2010; Rossi et al. 2012), and no development of resistance against the botanicals has yet been reported (Sharma et al. 1992). In some previous investigations, the effect of the essential oil and extracts of the *Mentha* species of pests was reported (Regnault-Roger and Hamraoui 1994; Pascual-villalobos and Robledo 1998; Mahmoudvand et al. 2011; Michaelakis et al. 2011) especially against the mosquito's species (Sukumar et al. 1991; Ansari et al. 2000; Traboulsi et al. 2002; Amer and Mehlhorn 2006).

The efficacy of the essential oils derived from the three *Mentha* species, *M. pulegium*, *M. piperita* and *M. spicata* against *C. pipiens* larvae, revealed that the *M. pulegium* and *M. piperita* oils were the most toxic ( $LC_{50}$ : 46.97 and 40.28ppm, respectively) and Pulegone was the most effective (27.23mg/l) among the major ingredients (Michaelakis et al. 2011).

According to the same author, Pulegone is 5-fold more active than Piperitone, this differentiation probably stands on the account of the location of the C-C double bond that these two molecules contain. We assume that in Pulegone the C-C double bond position on the chain group (isopropylidene versus isopropyl group) enhances the toxicity. Thus, Cetin et al. (2006) concluded that there was a high to the low lethal effect of the extracts of *Mentha pulegium* against the mosquito larvae, with  $LC_{50}$  values of 81.0ppm and Koliopoulos et al. (2010) confirmed the toxicity of *Mentha* ssp on the Culicidae larvae (*C. pipiens*).

The petroleum ether extract of *Moringa Oleifera* exhibited higher activity compared to the other extracts with  $LC_{50}$  values of 9.22ppm and 15.82ppm against *A. Gambiae* and *A. aegypti*, respectively (Nkya et al. 2014). Sayah et al. (2014) shows that the essential oil of *Citrus aurantium* (bitter orange), *Citrus sinensis* (orange) and *Pistacia lentiscus* presented an interesting larvicidal activity against *C. pipiens* ( $LC_{50}$ = 35, 64 and 62ppm and  $LC_{90}$ = 70, 120 and 160 ppm, respectively).

Eight out of the 22 essential oils (aniseed, calamus, cinnamon, clove, lemon, orange, thyme, and tulsi oils) gave promising results on the larvicidal activity against *C. quinquefasciatus* and the clove oil was found to be the most toxic to the larvae (Ramar et al. 2013). Cinnamon (*Cinnamomum cassia*), Sabina (*Sabina vulgaris*), White camphor (*Cinnamomum glanduliferum*), and Wintergreen (*Lix chinensis*) oils exhibited a larvicidal activity against the dengue vector *Aedes aegypti* larvae (Shaalan 2009). Prajapathi et al. (2005) found that the essential oils of the various parts of the 10 medicinal plants showed the different mosquitocidal activity against *A. stephensi*, *A. aegypti* and *C. quinquefasciatus*. The phytoproducts possess different bioactive components that can be used as general toxicants against the various larval stages of mosquitoes (Shaalan et al. 2005).

### 4. CONCLUSION

In the present study, it can be concluded that the essential oil of *M. pulegium* with pulegone, and eucalyptol as the major constituents, was found to exhibit a potent larvicidal activity against the *C. pipiens* and *A. caspius* larvae. Moreover, the *M. pulegium* essential oil appeared to be more toxic against the *A. caspius* larvae in comparison to the *C. pipiens* larvae. The essential oils can greatly contribute to reducing both the environmental chemicalisation and the population density of the mosquitoes.

### Acknowledgements

This work was supported by the National Fund for Scientific Research to Pr N Soltani (Laboratory of Applied Animal Biology) and the Ministry of High Education and Scientific Research of Algeria (CNEPRU Project to Pr F Tine-Djebbar).

### Conflict of Interest

Pr Tine-Djebbar F has received research grants from Ministry of High Education and Scientific Research of Algeria (CNEPRU project, grant number **D01N01UN120120130005**). Pr. Soltani N is a headmaster of Laboratory of Applied Animal Biology, University of Badji mokhtar, Annaba Guenez R declares that she has no conflict of interest.

**Author's Contribution:** All authors are equally contributed to this study.

Guenez Radja, Tine-Djebbar Fouzia and Soltani Nouredine, designed and carried out the experimental study. Tine-Djebbar Fouzia analysed data. Tine-Djebbar Fouzia, Guenez Radja and


Soltani Nouredine wrote the manuscript. All authors approved the manuscript.

#### REFERENCES

- Abbott W.S (1925) A method of computing the effectiveness of an insecticide. *Econ Entomol* 18: 265-267.
- Abdullah I.H (2009) Characterization and biological activities of essential oils of some species of Lamiaceae. Ph.D. Thesis, Agricultural Univ. Faisalabad, Pakistan, unpublished.
- Aberchane M, Fechtal M, Chaouch A, Bouayoune T (2001) Effect of time and distillation techniques on essential oils yield and quality of atlas cedar (*Cedrus atlantica* M.). *Ann Rech Forest Maroc* 34: 110-118.
- Amalich S, Zerkani H, Cherrat A, Dédianhoua N, Soro K, Bourakhouadar M, Mahjoubi M, EL Hilali F, Zair T (2016) Study on *Mentha pulegium* L. from M'rirt (Morocco): Antibacterial and antifungal activities of a pulegone-rich essential oil. *J Chem Pharm Res* 8(5): 363-370.
- Ambindei W.A, Dongmo Jazet P.M, Tatsadjieu L.N, Priya P, Manilal V.B, Krishnakumar B, Amvam Zollo P.H (2017) Effect of the essential oils of *Thymus vulgaris*, *Cinnamomum zeylanicum* and *Mentha piperita* on fungal growth and morphology. *Afr J Biotech* 16(9): 388-399.
- Amer A, Mehlhorn H (2006) Larvicidal effects of various essential oils against *Aedes*, *Anopheles*, and *Culex* larvae (Diptera, Culicidae). *Parasitol Res* 99: 466-472.
- Ansari M.A, Razdan R.K (1995) Relative efficacy of various oils in repelling mosquitoes. *Ind J Malar* 32: 104-111
- Ansari M.A, Vasudevan P, Tandon M, Razdan R.K (2000) Larvicidal and mosquito repellent action of peppermint (*Mentha piperita*) oil. *Biores Technol* 71: 267-271.
- Anonym (1983) Informal consultation on insect growth regulators. WHO/VBC/83.
- Benayad N (2008) Les huiles essentielles extraites des plantes médicinales marocaines : moyen efficace de lutte contre les ravageurs des denrées alimentaires stockées. Rapport d'étude. Faculté des sciences de Rabat, Maroc, 63p.
- Ben Slimane B, Ezzine O, Dhahri S, Chograni H, Ben Jamaa M.L (2015) Chemical composition of *Rosmarinus* and *Lavandula* essential oils and their insecticidal effects on *Orgyia trigotephras*. *J Coast Life Med* 3(1): 64-69.
- Bhatnagar M, Kapur K.K, Jalers S, Sharma S.K (2005) Laboratory evaluation of insecticidal properties of *Ocimum basilicum* L., *Ocimum sanctum* L. plants essential oils, and their major constituents against vector mosquito species. *J Entomol Res* 17: 21-26.
- Boudjelida H, Bouaziz A, Soin T, Smaghe G, Soltani N (2005) Effects of ecdysone agonist halofenozide against *Culex pipiens*. *Pest Biochem Physiol* 83(2/3): 115-123.
- Boughdad A, Elkasimi R, Kharchafi M (2011) Activité insecticide des huiles essentielles de *Mentha* sur *Callosobrochus maculatus* (F) (Coleoptera, Bruchidea). AFPP, 9ème Conférence Internationale sur les ravageurs en Agriculture, 2011 Octobre 26-27, Montpellier, France. 9p.
- Bouguerra N, Tine-Djebbar F, Soltani N (2017) Algerian *Thymus vulgaris* essential oil: chemical composition and larvicidal activity against the mosquito *Culex pipiens*. *Inter J Mosq Res* 4(1): 37-42.
- Boukhebt H, Chaker AN, Belhadj H, Sahli F, Ramdhani M, Laouer H, Harzallah D (2011) Chemical composition and antibacterial activity of *Mentha pulegium* L. and *Mentha spicata* L. essential oils. *Schol Res Libr* 3(4): 267-275.
- Bourkhiss M, Hnach M, Bourkhiss B, Ouhssine M, Chaouch A, Satrani B (2009) Effet de séchage sur la teneur et la composition chimique des huiles essentielles de *Tetraclinis articulata* (Vahl) Masters. *Agrosolutions* 20(1): 29-44.
- Bourkhiss M, Hnach M, Lakhlifi T, Boughdad A, Farah A, Satrani B (2011) Effect of age and vegetative stage on essential oil content and chemical composition of *Thuya articulata*. *Les Technologies de Laboratoire* 6(23): 64-68.
- Brunhes J, Rhaim A, Geoffroy B, Hervy J.P (2000) Mosquitoes of the Mediterranean Africa: Software Identification and Education, CD-ROM (Didactiques), IRD, Paris, France; IPT, Tunis, Tunisia.
- Cetin H, Cinbilgel I, Yanikoglu A, Gokceoglu M (2006) Larvicidal activity of some Labiatae (Lamiaceae) plant extracts from Turkey. *Phytoth Res* 20(12): 1088-1090.
- Cherrat L, Espina L, Bakkali M, Pagán R, Laglaoui A (2014) Chemical composition, antioxidant and antimicrobial properties of *Mentha pulegium*, *Lavandula stoechas* and *Satureja calamintha* Scheele essential oils and an evaluation of their bactericidal effect in combined processes. *Innov Food Sci Emerg Technol* 22: 221-229.
- Choochate W, Chaiyasit D, Kanjanapothi D, Rattanachanpichai E, Jitpakdi A, Tuetun B (2005) Chemical composition and antimosquito potential of rhizome extract and volatile oil derived from *Curcuma aromatica* against *Aedes aegypti* (Diptera: Culicidae). *J Vect Ecol* 30: 302-309.
- Dahchar Z, Bendali-Saoudi F, Soltani N (2016) Larvicidal activity of some plant extracts against two mosquito species *Culex pipiens* and *Culiseta longiareolata*. *J Entomol Zool Stud* 4(4): 346-350.
- Derwich E, Benziane Z, Boukir A (2009) Chemical compositions and insecticidal activity of essential oils of three plants *Artemisia* sp: *Artemisia herba-alba*, *Artemisia absinthium* and *Artemisia Pontica* (Morocco). *Electr J Envir Agr Food Chem* 8(11): 1202-1211.
- Dharmagadda V.S.S, Naik S.N, Mittal P.K, Vasudevan P (2005) Larvicidal activity of *Tagetes patula* essential oil against three mosquito species. *Biores Technol* 96: 1235-1240.
- El Akhal F, Greche H, Ouazzani CF, Guemmouh R, El Ouali Lalami A (2015) Chemical composition and larvicidal activity of *Culex pipiens* essential oil of *Thymus vulgaris* grown in Morocco. *J Mater Environ Sci* 1: 214-219.
- El Akhal F, Guemmouh R, Maniar S, Taghzouti K, El Ouali Lalami A (2016) Larvicidal activity of essential oils of *Thymus vulgaris* and *Origanum majorana* (Lamiaceae) against of the malaria vector *Anopheles labranchiae* (Diptera: Culicidae). *Inter J Pharm Pharmaceutical Sci* 8(3): 372-376.
- El-Ghorab A.H (2006) The chemical composition of *Mentha pulegium* L. essential oil from Egypt and its antioxidant activity. *J Ess Oil Bear Plant* 9: 183-195.

29. El Ouali Lalami A, EL-Akhal F, Maniar S, Ezzoubi Y, Taghzouti K (2016) Chemical constituents and larvicidal activity of essential oil of *Lavandula stoechas* (Lamiaceae) from morocco against the malaria vector *Anopheles labranchiae* (Diptera: Culicidae). *Inter. J Pharm Phytochem Res* 8(3): 505-511.
30. Finney D.J (1971) *Probit analysis*. 3rd ed., Cambridge University Press, London and New York.
31. Ghosh A, Chowdhury N, Chandra G (2012) Plant extracts as potential mosquito larvicides. *Ind J Med Res* 135: 581-598.
32. Govindarajan M (2010) Larvicidal and repellent activities of *Sida acuta* Burm. F. (Family: Malvaceae) against three important vector mosquitoes. *Asian Pacific J Trop Med* 3(9): 691-695.
33. Germplasm Resources Information Network (2010) (Online Database) Beltsville (MD): National Germplasm Resources Laboratory.
34. Hafedh H, Najla T, Emira N, Mejdi S, Hanen F, Riadh K, Amina B (2009) Biological activities of the essential oils and methanol extract of two cultivated mint species (*Mentha longifolia* and *Mentha pulegium*) used in the Tunisian folkloric medicine. *W J Microbiol Biotech* 25: 2227-2238.
35. Hilan C, Sfeir R, El Hage R, Jawich D, Frem M.E, Jawhar K (2007) Evaluation of the antibacterial activities of *Ferula hermonis* (Boiss.). *Leb Sci J* 8(2): 135-151.
36. Hmiri S, Rahouti M, Zakaria H, Satrani B, Ghanmi M, El Ajjour M (2010) Évaluation du potentiel antifongique des huiles essentielles de *Mentha pulegium* et d'*Eucalyptus camaldulensis* dans la lutte biologique contre les champignons responsables de la détérioration des pommes en conservation. *Bul Soc Roy Sci Liège* 80: 824-836.
37. Himmi O, Dakki M, Trari B, El Agbani M.A, 1995 *The Culicidae of Morocco: Identification Keys with Biological and Ecological Data (Work of the Scientific Institute), Zoology of Series*, Rabat, Morocco
38. Imelouane B, Amhamdi H, Wathelet J.P, Ankit M, Khedid K, El Bachiri A (2009) Chemical Composition and Antimicrobial Activity of Essential Oil of Thyme (*Thymus vulgaris*) from Eastern Morocco. *Inter J Agr Biol* 11(2): 205-208.
39. Kastner G (1969) The dependance of field and quality of thyme grown as a two year crop on the date of harvesting and the cutting height in the autumn of the year of sowing. *Pharmazie* 24: 226-235.
40. Koliopoulos G, Pitarokili D, Kioulos E, Michaelakis A, Tzakou O (2010) Chemical composition and larvicidal evaluation of *Mentha*, *Salvia*, and *Melissa* essential oils against the West Nile virus mosquito *Culex pipiens*. *Parasitol Res* 107: 327-335.
41. Kokkini S, Karousou R, Hanlidou E, Lanaras T (2004) Essential oil composition of Greek (*Origanum vulgare* ssp *hirtum*) and Turkish (*O. onites*) oregano: A tool for their distinction. *J Ess Oil Res* 16 (4): 334-338.
42. Koul O, Walia S, Dhaliwal G.S (2008) Essential oils as green pesticides: potential and constraints. *Biopest Inter* 4(1): 63-84.
43. Laggoune S, Öztürk M, Erol E, Duru M.E, Abaza I, Kabouche A, Kabouche Z (2016) Chemical composition, antioxidant and antibacterial activities of the essential oil of *Mentha spicata* L. from Algeria. *J Mater Envir Sci* 7(11): 4205-4213.
44. Lorenzo D, Paz D, Eduardo D, Philip D, Roser Villa D, Cañigual S (2002) Essential oils of *Mentha pulegium* and *Mentha rotundifolia* From Uruguay. *Braz Arch Biol Technol* 45: 519-524.
45. Mahboubi M, Haghi G (2008) Antimicrobial activity and chemical composition of *Mentha pulegium* L. essential oil. *J Ethnopharm* 119(2): 325-327.
46. Mahmoudvand M, Abbasipour H, Basij M, Hosseinpour M.H, Rastegar F, Nasiri M.B (2011) Fumigant toxicity of some essential oils on adults of some stored-product pests. *Chil J Agr Res* 71(1): 83-89
47. Michaelakis A, Papachristos D, Kimbaris A, Polissiou M (2011) Larvicidal evaluation of three *Mentha* species essential oils and their isolated major components against the West Nile virus mosquito. *Hell Plant Protec J* 4: 35-43.
48. Mkaddem M, Bousaid M, Ben Fadhel N (2007) Variability of volatiles in 9Tunisian *Mentha pulegium* L. (Lamiaceae). *J Ess Oil Res* 19: 211-215.
49. Nkya J.P, Erasto P, Chacha M (2014) Larvicidal against mosquito vectors and brine shrimp activities of extracts from the flowers of *Moringa oleifera* Lam. *Amer J Res Com* 2(8): 17-18.
50. Okoh O.O, Sadimenko A.A, Afolayan A.J (2007) The effects of age on the yield and composition of the essential oils of *Calendula officinalis*. *J Appl Sci* 7(23): 3806-3810.
51. Onyambu G.K, Maranga R, Ndungu M, Kareru G, Wanjoya A (2015) GC-MS analysis of pesticidal essential oils from Kenyan plants. *Afr J Biotech* 14 (13): 1158-1166.
52. Ouakouak H, Chohra M, Denane M (2015) Chemical composition, antioxidant activities of the essential oil of *Mentha pulegium* L, South East of Algeria. *Inter Let Nat Sci* 39: 49-55.
53. Ouraini D, Agouni A, Alaoui M.I, Alaoui K (2005) Therapeutic approach of dermatophytes by HE Moroccan herbs. *Herb Med Phytoth* 1: 3-12.
54. Pascual-villalobos M.S, Robledo A (1998) Screening for anti-insect activity in Mediterranean plants. *J Indust Crops Prod* 8(3): 115-120.
55. Pavela R (2008a) Insecticidal properties of several essential oils on the house fly (*Musca domestica* L.). *Phytoth Res* 22: 274-278.
56. Pavela R (2008b) Larvicidal effects of various Euro-Asiatic plants against *Culex quinquefasciatus* Say. larvae (Diptera: Culicidae). *Parasitol Res* 102: 555-559.
57. Pavela R, Kaffková K, Kumšta M (2014) Chemical composition and larvicidal activity of essential oils from different *Mentha* and *Pulegium* species against *Culex quinquefasciatus* Say (Diptera: Culicidae) *Plant Protec Sci* 50 (1): 36-42.
58. Prajapathi N.D, Prajapathi T, Jaipura S (2005) Advances in medicinal plants. *Asian medicinal plant and health care trust*, Jodhpur 1.
59. Ramar M, Gabriel Paulraj M, Ignacimuthu S (2013) Preliminary screening of plant essential oils against larvae of *Culex quinquefasciatus* Say (Diptera: Culicidae). *Afr J Biotech* 12(46): 6480-6483.

60. Regnault-Roger C, Hamraoui A (1994) Reproductive inhibition of *Acanthoscelides obtectus* Say (Coleoptera), bruchid of kidney bean (*Phaseolus vulgaris* L.) by some aromatic essential oils. *Crop Protec* 13:624-28.
61. Rocha D, Novo M, Matos O.C, Figueiredo A, Delgado M.D, Cabral M, Liberato M, Moiteiro C (2015) Potential of *Mentha pulegium* for mosquito control. *Revista de Ciências Agrárias* 38(2): 155-165.
62. Rossi E, Cosimi S, Loni A (2012) Bioactivity of essential oils from mediterranean plants: insecticidal properties on *Sitophilus zeamais* and effects on seed germination. *J Entomol* 9: 403-412.
63. Sayah M.Y, El Ouali lalami A.H, Greech H, Errachidi F, Rodi El Kandri Y, Ouazzani Chahdi F (2014) Larvicidal activity of aromatic plant extracts on larvae of mosquitoes vectors of parasitic diseases. *Inter. J Innov Appl Stud* 7(3): 832-842.
64. Shaalan E.A (2009) Larvicidal activity of some volatile oils against the dengue vector *Aedes aegypti*. 45 Annual Scientific Seminar of Malaysian Society of Parasitology and Tropical Medicine "Impact of animal hosts on disease transmission and public health" 18-19 March. Kuala Lumpur, Malaysia.
65. Sardashti A, Adhami Y (2013) Chemical composition of the essential oil of *Mentha pulegium* L. from Taftan Area by means of gas chromatography/mass spectrometry (GC/MS). *J Med Plant Res* 7(40): 3003-3007.
66. Samarasekera R, Weerasinghe I.S, Hemalal K.D.P (2008) Insecticidal activity of menthol derivatives against mosquitoes. *Pest Manag Sci* 64: 290-295.
67. Schaffner F, Angel G, Geoffroy B, Hevry J.P, Rhaïem A, Brunhes J (2001) Moustiques d'Europe. Institut de recherche pour le développement IRD. Logiciel d'identification.
68. Sharma R.N, Gupta A.S, Patwardhan S.A, Hebbal-ker D.S, Tare V, Bhonde S.B (1992) Bioactivity of Lamiaceae plants against insects. *Ind J Exp Biol* 30: 244-246
69. Shaalan E.A.S, Canyonb D, Younesc M.W.F, Abdel-Wahaba H, Mansoura A.H (2005) A review of botanical phytochemicals with mosquitocidal potential. *Envir Inter* 3: 1149-1166.
70. Silva L.F, Cardoso M.G, Batista L.R, Gomes M.S, Rodrigues L.M.A, Rezende D.A.C.S, Teixeira M.L, Carvalho M.S.S, Santiago J.A, Nelson D.L (2015) Chemical characterization, antibacterial and antioxidant activities of essential oils of *Mentha viridis* L. and *Mentha pulegium* L. (L). *Amer J Plant Sci* 6: 666-675.
71. Skaltsa H.D, Demetzos C, Lazari D, Sokovic M (2003) Essential oil analysis and antimicrobial activity of eight *Stachys* species from Greece. *Phytochem* 64: 743-752.
72. Soković M.D, Vukojević J, Marin P.D, Brkić D.D, Vajs V, Van Griensven L.J.L.D (2009) Chemical composition of essential oils of *Thymus* and *Mentha* species and their antifungal activities. *Molecules* 14(1): 238-249.
73. Stoyanova A, Georgie V, Kula J, Majda T (2005) Chemical composition of the essential oil of *Mentha pulegium* from Bulgaria. *J Ess Oil Res* 17: 475-477.
74. Sukumar K, Perich M.J, Boobar L.R (1991) Botanical derivatives in mosquito control: a review. *J Amer Mosq Contr Assoc* 7: 210-237.
75. Swaroop S, Gilroy A.B, Uemura K (1966) Statistical methods in malaria eradication. Geneva: World Health Organisation.
76. Traboulsi A.F, Taoubi K, El-Haj S, Bessiere J.M, Rammal S (2002) Insecticidal properties of essential plant oils against the mosquito *Culex pipiens molestus* (Diptera: Culicidae). *Pest Manag Sci* 58(5): 491-5.
77. Tine-Djebbar F, Bouabida H, Soltani N (2016) Répartition spatio-temporelle des Culicidés dans la région de Tébessa. Editions universitaires européennes ISSN/ISBN : 978-3-639-50856-7.
78. Urzúa A, Santander R, Echeverría J, Cabezas N, Palacios S.M, Rossi Y (2010) Insecticide properties of the essential oils from *Haplopappus foliosus* and *Bahia ambrosoides* against the house fly, *Musca domestica* L. *J Chil Chem Soc* 55: 392-395.
79. Verma R.S, Rahman L, Verma R.K, Chauhan A, Yadav A.K, Singh A (2010) Essential oil composition of menthol mint (*Mentha arvensis*) and peppermint (*Mentha piperita*) cultivate at different stages of plant growth from Kumaon region of western Himalaya. *Inter. J Med Arom Plants* 1: 13-18.
80. Vernet P.H, Guillerm J.L, Gouyon P.H (1977) Le polymorphisme chimique de *Thymus vulgaris* L. (Labiée), I. Répartition des formes chimiques en relation avec certains facteurs écologiques. *Ecologica Plantarum* 128(2): 159-179.
81. Vian M.A, Fernandez X, Visinoni F, Chemat F (2008) Microwave hydrodiffusion and gravity, a new technique for extraction of essential oils. *J Chrom A* 1190: 14-17.
82. Zantar S, El Garrouj D, Pagán R, Chabi M, Laglaoui A, Bakkali M, Zerrouk M.H (2015) Effect of harvest time on yield, chemical composition, antimicrobial and antioxidant activities of *Thymus vulgaris* and *Mentha pulegium* Essential Oils. *Europ J Med Plants* 8(2): 69-77.



**Fig. 1.** Typical GC-MS Chromatograms of *M. pulegium* essential oils showing the separation of chemical components using HP-5MS column (Area as a function of the time in min)

**Table 1.** Chemical composition of *M. pulegium* oils: retention time (RT) and concentration (%) of different constituents.

Pic	Compound	RT	%
<b>MONOTERPENES HYDROCARBONS</b>			
1	$\alpha$ - Pinene	6.33	0.26
2	$\beta$ - Phellandrene	7.44	0.38
3	3-Carene Bicyclo (3.1.0) Hexane.4-Methyl-1-(1-Methylethyl) Didehydro Deriv	7.55	0.32
5	L- $\beta$ Pinene	11.90	1.90
13	D-Limonene	22.33	0.63
<b>OXYGENATED HYDROCARBONS</b>			
4	Eucalyptol	9.38	10.44
6	P-Menthone 2-Ethyl-5-Propyle	14.09	5.46
7	2- Ethyl-5-Propylcyclo Pentanone	14.53	3.78
8	Menthol	14.87	0.68
9	Isopulegone	14.99	1.12
10	$\alpha$ - Terpeneol	15.64	0.63
11	Pulegone	17.87	72.50
12	Piperitenone	21.99	1.51
14	Eugenol Methyl Ether	24.62	0.37
<b>Total (%)</b>			<b>99.29</b>

**Table 2.** Efficacy of essential oil of *Mentha pulegium* applied on fourth instar larvae of *Culex pipiens* and *Aedes caspius*: corrected mortality ( $m \pm SD$ , n = 4 replicates each containing 25 larvae).

Species	12.5 ppm	25 ppm	50 ppm	75 ppm	100 ppm	P value
<i>C. pipiens</i>	5.38 $\pm$ 2.26	24.27 $\pm$ 3.59	54.00 $\pm$ 5.29	66.22 $\pm$ 6.01	95.94 $\pm$ 4.00	p $\leq$ 0.001
<i>A. caspius</i>	16.33 $\pm$ 6.63	28.77 $\pm$ 0.67	57.55 $\pm$ 1.34	91.83 $\pm$ 3.91	100 $\pm$ 0.00	p $\leq$ 0.001

**Table 3.** Larvicidal activity of *Mentha pulegium* oils against fourth instar of *Aedes caspius* and *Culex pipiens* after 24 hours of treatment

Species	Regression equation	LC <sub>50</sub>		LC <sub>90</sub>		Slope
		ppm	95% FL	ppm	95% FL	
<i>C. pipiens</i>	y = 3.706 x - 0.886	38.75	(35.71 - 42.04)	85.91	(70.71 - 104.37)	1.85
<i>A. caspius</i>	y = 3.542 x - 0.534	28.16	(25.86 - 30.65)	53.75	(43.37 - 66.61)	1.65